

A transience condition for a class of one-dimensional symmetric Lévy processes

Nikola Sandrić
Department of Mathematics
Faculty of Civil Engineering, University of Zagreb, Zagreb, Croatia
Email: nsandric@grad.hr

August 22, 2013

Abstract

In this paper, we give a sufficient condition for the transience for a class of one-dimensional symmetric Lévy processes. More precisely, we prove that a one-dimensional symmetric Lévy process with the Lévy measure $\nu(dy) = f(y)dy$ or $\nu(\{n\}) = p_n$, where the density function $f(y)$ is such that $f(y) > 0$ a.e. and the sequence $\{p_n\}_{n \geq 1}$ is such that $p_n > 0$ for all $n \geq 1$, is transient if

$$\int_1^\infty \frac{dy}{y^3 f(y)} < \infty \quad \text{or} \quad \sum_{n=1}^\infty \frac{1}{n^3 p_n} < \infty.$$

Similarly, we derive an analogous transience condition for one-dimensional symmetric random walks with continuous and discrete jumps.

Keywords and phrases: characteristics of a semimartingale, electrical network, Lévy measure, Lévy process, random walk, recurrence, transience

1 Introduction

Let $(\Omega, \mathcal{F}, \mathbb{P})$ be a probability space and let $\{L_t\}_{t \geq 0}$ be a stochastic process on $(\Omega, \mathcal{F}, \mathbb{P})$ taking values in \mathbb{R}^d , $d \geq 1$. The process $\{L_t\}_{t \geq 0}$ is called a *Lévy process* if $L_0 = 0$ \mathbb{P} -a.s., if it has stationary and independent increments and if it has càdlàg paths \mathbb{P} -a.s. (that is, if its trajectories are right-continuous with left limits \mathbb{P} -a.s.). Having these properties, every Lévy process can be completely and uniquely characterized through the characteristic function of a single random variable L_t , $t > 0$, that is, by the famous *Lévy-Khintchine formula* we have

$$\mathbb{E}[\exp\{i\langle \xi, L_t \rangle\}] = \exp\{-t\psi(\xi)\}, \quad t \geq 0,$$

where

$$\psi(\xi) = i\langle \xi, b \rangle + \frac{1}{2}\langle \xi, c\xi \rangle + \int_{\mathbb{R}^d} (1 - \exp\{i\langle \xi, y \rangle\} + i\langle \xi, y \rangle 1_{\{|y| \leq 1\}}(y)) \nu(dy).$$

Here b is a vector in \mathbb{R}^d , c is a symmetric nonnegative-definite $d \times d$ matrix and $\nu(dy)$ is a σ -finite Borel measure on \mathbb{R}^d satisfying

$$\nu(\{0\}) = 0 \quad \text{and} \quad \int_{\mathbb{R}^d} \min\{1, y^2\} \nu(dy) < \infty.$$

The measure $\nu(dy)$, the triplet (b, c, ν) and the function $\psi(\xi)$ are called the *Lévy measure*, the *Lévy triplet* and the *characteristic exponent* of the Lévy process $\{L_t\}_{t \geq 0}$, respectively. Further, recall that the vector b , the matrix c and the Lévy measure $\nu(dy)$ correspond to the deterministic part (shift), the continuous (Brownian) part and the jumping part of the Lévy process $\{L_t\}_{t \geq 0}$, respectively.

In this paper, we consider the transience and recurrence property of Lévy processes. A Lévy process $\{L_t\}_{t \geq 0}$ is said to be *transient* if

$$\mathbb{P}\left(\lim_{t \rightarrow \infty} |L_t| = \infty\right) = 1,$$

and *recurrent* if

$$\mathbb{P}\left(\liminf_{t \rightarrow \infty} |L_t| = 0\right) = 1.$$

It is well known that every Lévy process is either transient or recurrent (see [Sat99, Theorem 35.3]). An equivalent definition (characterization) of the transience and recurrence property of Lévy processes can be given through the sojourn times. A Lévy process $\{L_t\}_{t \geq 0}$ is transient if and only if

$$\mathbb{E}\left[\int_0^\infty 1_{\{|L_t| < a\}}(L_t) dt\right] < \infty \quad \text{for all } a > 0.$$

Similarly, a Lévy process $\{L_t\}_{t \geq 0}$ is recurrent if and only if

$$\mathbb{E}\left[\int_0^\infty 1_{\{|L_t| < a\}}(L_t) dt\right] = \infty \quad \text{for all } a > 0$$

(see [Sat99, Theorem 35.4]). The above characterizations of the transience and recurrence property are not applicable in most cases. A more operable characterization, by using the nice analytical characterization of Lévy processes through the Lévy-Khintchine formula, has been given by the well-known *Chung-Fuchs criterion*. A Lévy process $\{L_t\}_{t \geq 0}$ is transient if and only if

$$\int_{\{|\xi| < a\}} \operatorname{Re}\left(\frac{1}{\psi(\xi)}\right) d\xi < \infty \quad \text{for some } a > 0$$

(see [Sat99, Corollary 37.6 and Remark 37.7]). Again, in many situations this criterion is also not applicable. More precisely, for a given Lévy triplet (b, c, ν) it is not always easy to compute the integral part of the characteristic exponent as well as the integral appearing in the Chung-Fuchs criterion. According to this, the aim of this paper is to derive a sufficient condition for the transience for Lévy processes in terms of the Lévy triplet. Let us remark that analogous definitions and characterizations of the transience and recurrence property hold also for random walks (see [Dur10, Chapter 4]). Recall that a *random walk* is a stochastic process $\{S_n\}_{n \geq 0}$ defined on a probability space $(\Omega, \mathcal{F}, \mathbb{P})$ taking values in \mathbb{R}^d , $d \geq 1$, defined by $S_0 := 0$ and $S_n := \sum_{i=1}^n J_i$, where $\{J_n\}_{n \geq 1}$ is a sequence of i.i.d. random variables called the *jumps* of $\{S_n\}_{n \geq 0}$.

As already mentioned, in this paper we consider the one-dimensional symmetric case only. Note that, according to [Sat99, Theorem 37.8] and [Dur10, Theorem 4.2.13], the limitation to the one-dimensional case is not too big restriction since it is well known that every d -dimensional, $d \geq 3$, Lévy process and random walk are transient. Further, recall that a stochastic process $\{X_t\}_{t \in \mathbb{T}}$ is *symmetric* if $\{X_t\}_{t \in \mathbb{T}} \stackrel{d}{=} \{-X_t\}_{t \in \mathbb{T}}$, where $\mathbb{T} = [0, \infty)$ or $\{0, 1, 2, \dots\}$ and $\stackrel{d}{=}$ means that the processes $\{X_t\}_{t \in \mathbb{T}}$ and $\{-X_t\}_{t \in \mathbb{T}}$ have the same finite-dimensional distributions. In the Lévy process case, $\{L_t\}_{t \geq 0}$ is symmetric if and only if $b = 0$ and $\nu(dy)$ is a symmetric measure, that is, $\nu(B) = \nu(-B)$ holds for all Borel sets $B \subseteq \mathbb{R}^d$ (see [Sat99, Exercise 18.1]), while a random walk is symmetric if and only if its jumps have a symmetric distribution. Now, let us state the main results of this paper.

Theorem 1.1. Let $\{L_t\}_{t \geq 0}$ be a one-dimensional symmetric Lévy process with the Lévy measure $\nu(dy) = f(y)dy$ or $\nu(\{n\}) = p_n$, where the density function $f(y)$ is such that $f(y) > 0$ a.e. and the sequence $\{p_n\}_{n \geq 1}$ is such that $p_n > 0$ for all $n \geq 1$. Then, $\{L_t\}_{t \geq 0}$ is transient if

$$\int_1^\infty \frac{dy}{y^3 f(y)} < \infty \quad \text{or} \quad \sum_{n=1}^\infty \frac{1}{n^3 p_n} < \infty. \quad (1.1)$$

By using [Sat99, Theorem 38.2], the above transience condition can be generalized to the general symmetric case.

Theorem 1.2. Let $\{L_t^1\}_{t \geq 0}$ and $\{L_t^2\}_{t \geq 0}$ be one-dimensional symmetric Lévy processes with the Lévy measures $\nu_1(dy)$ and $\nu_2(dy)$. Further, let $\nu_1(dy)$ be as in Theorem 1.1 and let it satisfy the condition (1.1). If

$$\int_0^\infty y^2 |\nu_1 - \nu_2|(dy) < \infty,$$

then the transience property of $\{L_t^1\}_{t \geq 0}$ implies the transience property of $\{L_t^2\}_{t \geq 0}$. Here $|\cdot|$ denotes the total variation norm on the space of signed measures.

Let us remark that the same transience condition holds also in the case of one-dimensional symmetric random walks. More precisely, let $\{S_n\}_{n \geq 0}$ be a one-dimensional symmetric random walk with jumps $\mathbb{P}(J_1 \in dy) = f(y)dy$ or $\mathbb{P}(J_1 = n) = p_n$, where $f(y) > 0$ a.e. and $p_n > 0$ for all $n \geq 1$, then the condition (1.1) implies the transience property of $\{S_n\}_{n \geq 0}$. Also, let us remark that, according to [Sat99, Theorem 38.2] or [She62, Lemma 1.2], the assumptions $f(y) > 0$ a.e. and $p_n > 0$ for all $n \geq 1$ can be relaxed. More precisely, it suffices to demand positivity of $f(y)$ and p_n on the complement of a compact set.

A simple application of the condition (1.1) is in the class of stable processes. Recall that a one-dimensional symmetric stable Lévy process or a random walk is given by the characteristic exponent $\psi(\xi) = \gamma|\xi|^\alpha$ or by the Lévy triplet $(0, c, \nu)$, where $\gamma > 0$, $\alpha \in (0, 2]$,

$$c = \begin{cases} 0, & \alpha < 2 \\ 2\gamma, & \alpha = 2 \end{cases} \quad \text{and} \quad \nu(dy) = \begin{cases} \gamma \frac{\alpha 2^{\alpha-1} \Gamma(\frac{\alpha+1}{2})}{\pi^{\frac{1}{2}} \Gamma(1-\frac{\alpha}{2})} |y|^{-\alpha-1} dy, & \alpha < 2 \\ 0, & \alpha = 2. \end{cases}$$

It is well known, as a consequence of the Chung-Fuchs criterion, that this process is transient if and only if $\alpha < 1$. Further, recall that a probability density function of a one-dimensional symmetric stable distribution behaves like $c_\alpha |y|^{-\alpha-1}$ when $|y| \rightarrow \infty$, for $\alpha \in (0, 2)$ and

$$c_\alpha = \begin{cases} \frac{\gamma}{2}, & \alpha = 1 \\ \frac{\gamma}{\pi} \Gamma(\alpha + 1) \sin\left(\frac{\pi\alpha}{2}\right), & \alpha \neq 1 \end{cases}$$

(see [Sat99, Remark 14.18]). Now, as a simple consequence of Theorem 1.1, we get a new proof for the transience property of one-dimensional symmetric stable Lévy processes and random walks.

Corollary 1.3. A one-dimensional symmetric stable Lévy process or a random walk is transient if $\alpha < 1$.

Note that the above corollary implies that the function $y \mapsto y^3$, appearing in the condition (1.1), is optimal in the class of power functions.

The transience and recurrence property of one-dimensional symmetric Lévy processes in terms of the Lévy triplet has already been studied in the literature. Namely, in [Sat99, Theorem 38.3] (see also [She62, Theorem 5]) it has been proved that a one-dimensional symmetric Lévy process $\{L_t\}_{t \geq 0}$ with the Lévy measure $\nu(dy)$ is recurrent if

$$\int_1^\infty \left(\int_0^y z \nu(\max\{1, z\}, \infty) dz \right)^{-1} dy = \infty. \quad (1.2)$$

Intuitively, the condition (1.2) measures the speed of divergence of the second moment of $\nu(dy)$, and, regarding this speed, it concludes the recurrence property. Clearly, if $\nu(dy)$ has finite second moment, then $\{L_t\}_{t \geq 0}$ is recurrent. Thus, if the second moment of $\nu(dy)$ diverges slow enough, then $\{L_t\}_{t \geq 0}$ is recurrent. Similarly, the condition (1.1) measures the speed of divergence of the third moment of $\nu(dy)$. If the third moment of $\nu(dy)$ diverges fast enough, then $\{L_t\}_{t \geq 0}$ is transient.

Recall that a symmetric Borel measure $\rho(dy)$ on \mathbb{R} is *unimodal* if it is finite outside of any neighborhood of the origin and if $x \mapsto \rho(x, \infty)$ is a convex function on $(0, \infty)$. Equivalently, a symmetric Borel measure $\rho(dy)$ on \mathbb{R} is unimodal if it is of the form $\rho(dy) = a\delta_0(dy) + f(y)dy$, where $0 \leq a \leq \infty$ and the density function $f(y)$ is symmetric, decreasing on $(0, \infty)$ and it satisfies $\int_{|y| > \varepsilon} f(y)dy < \infty$ for all $\varepsilon > 0$ (see [Sat99, Chapter 5]). Note that measures with a discrete support are never unimodal. Now, if the Lévy measure $\nu(dy)$ is additionally unimodal, the condition (1.2) is also necessary for the recurrence property (see [Sat99, Theorem 38.3] or [She62, Theorem 5]). Also, note that unimodality of the $\nu(dy)$ and finiteness of (1.2) imply that $f(y) > 0$ a.e., and the condition (1.2) is stronger than the condition (1.1) (see Section 4 for the proof). Thus, the condition (1.1) is a generalization of the condition (1.2) in the case when the jumping measure is not unimodal. For the necessity of unimodality for the characterization of the transience property by the condition (1.2) see [Sat99, Theorems 38.2, 38.3 and 38.4 and Lemma 38.8] or [She62, Theorems 4 and 5] and [She64, Theorem 1].

Finally, we give an example where the condition (1.1) is more suitable than the Chung-Fuchs criterion and the condition (1.2). We consider an example of a Lévy process with “multiple indices of stability”. Let $\{L_t\}_{t \geq 0}$ be a one-dimensional symmetric Lévy process with the Lévy measure $\nu(\{n\}) = p_n$, where $p_{2n} = (2n)^{-\alpha-1}$ and $p_{2n-1} = (2n-1)^{-\beta-1}$ for $n \geq 1$ and $\alpha, \beta \in (0, \infty)$. For a continuous version of such process it suffices to interpolate the points $\{(i, p_i) : i \in \mathbb{Z}\}$. Now, clearly, if $\alpha < 1$ and $\beta < 1$, then the condition (1.1) implies the transience of $\{L_t\}_{t \geq 0}$. On the other hand, since $\nu(dy)$ is not unimodal, the condition (1.2) is not applicable, and the application of the Chung-Fuchs criterion leads to a non-trivial computation. Also, let us remark that the same example shows that the condition (1.1) is only sufficient for the transience. Indeed, assume that $\alpha < 1$ and $\beta \geq 1$. Then, since $\beta \geq 1$, the condition (1.1) fails to hold. On the other hand, since $\alpha < 1$, $\{L_t\}_{t \geq 0}$ is transient (see Section 4 for the proof).

Now, we explain our strategy of proving Theorem 1.1. The proof is divided in three steps. In the first step, by using electrical networks techniques, we prove Theorem 1.1 in the case of a random walk with discrete jumps. In the second step, we prove Theorem 1.1 in the case of a random walk $\{S_n\}_{n \geq 0}$ with continuous jumps $\mathbb{P}(J_1 \in dy) = f(y)dy$. More precisely, for $\delta > 0$ we define a discretization of $\{S_n\}_{n \geq 0}$ as a random walk $\{S_n^\delta\}_{n \geq 0}$ on $\delta\mathbb{Z}$ with the jump distribution $\mathbb{P}(J_1^\delta = \delta n) := \int_{\delta n - \frac{\delta}{2}}^{\delta n + \frac{\delta}{2}} f(y)dy$, $n \in \mathbb{Z}$. Next, by an “approximation approach” we prove that all the random walks $\{S_n^\delta\}_{n \geq 0}$, $\delta > 0$, are either transient or recurrent at the same time and their transience and recurrence property is equivalent with the transience and recurrence property of $\{S_n\}_{n \geq 0}$. Finally, by using the first step, we prove that the condition (1.1) for $\{S_n\}_{n \geq 0}$ implies the transience property of $\{S_n^1\}_{n \geq 0}$. And this accomplishes the proof of the second step. At the end, in the last step, we consider the case of Lévy processes. By using [Sat99, Theorem 38.2], it suffices to consider the situation of a compound Poisson process. Now, the proof follows from the first and second step. This accomplishes the proof of Theorem 1.1.

The paper is organized as follows. In Section 2, we give a proof of Theorem 1.1 for the case of discrete jumps. In Section 3, by using the results from Section 2, we proof Theorem 1.1 for the case of continuous jumps. Finally, in Section 4, we discuss some properties of the condition (1.1).

2 Discrete case

In this section, we prove the main step of the proof of Theorem 1.1. More precisely, we derive a sufficient condition for the transience for one-dimensional symmetric random walks on \mathbb{Z} .

Theorem 2.1. *Let $\{S_n\}_{n \geq 0}$ be a one-dimensional symmetric random walk on \mathbb{Z} with jumps $\mathbb{P}(J_1 = n) = p_n$, where the sequence $\{p_n\}_{n \geq 1}$ is such that $p_n > 0$ for all $n \geq 1$. Then the random walk $\{S_n\}_{n \geq 0}$ is transient if*

$$\sum_{n \geq 1} \frac{1}{n^3 p_n} < \infty. \quad (2.1)$$

Note that the same transience condition also holds in the case of a one-dimensional symmetric Lévy process $\{L_t\}_{t \geq 0}$ with a discrete supported Lévy measure $\nu(\{n\}) = p_n$, where $p_n > 0$ for all $n \geq 1$. Indeed, first note that

$$\{L_t\}_{t \geq 0} \stackrel{d}{=} \{S_{P_t}\}_{t \geq 0},$$

where $\{S_n\}_{n \geq 0}$ is a random walk with jumps $\mathbb{P}(J_1 = n) := \frac{1}{\nu(\mathbb{Z})} p_n$ and $\{P_t\}_{t \geq 0}$ is the Poisson process with parameter $\nu(\mathbb{Z})$ independent of $\{S_n\}_{n \geq 0}$. Now, the desired result follows from the definition of the transience in terms of sojourn times.

The proof of Theorem 2.1 is based on techniques and results from electrical networks. Let us introduce some notation we need. A *graph* is a pair $G = (V(G), E(G))$ where $V(G)$ is a set of *vertices* and $E(G)$ is a symmetric subset of $V(G) \times V(G)$, called the *edge set*. By symmetry we mean that $(u, v) \in E(G)$ if and only if $(v, u) \in E(G)$. For two vertices $u, v \in V(G)$ such that $(u, v) \in E(G)$, we say that u and v are *adjacent* and write $u \sim v$ and by e_{uv} we denote the edge which connects them. A *path* in a graph is a sequence of vertices where each successive pair of vertices is an edge in the graph. A graph is *connected* if there is a path from any of its vertices to any other. A *network* is a pair $N = (G, c)$, where G is a connected graph and c is a function $c : E(G) \rightarrow [0, \infty)$ called *conductance*. In the sequel we assume that a network N satisfies

$$c(u) := \sum_{v \sim u} c(e_{uv}) < \infty$$

for all $u \in V(G)$. A *random walk* on a network N is a time-homogeneous Markov chain $\{X_n\}_{n \geq 0}$ with the state space $V(G)$ and transition kernel

$$q_{uv} := \mathbb{P}(X_1 = v | X_0 = u) = \begin{cases} \frac{c(e_{uv})}{c(u)}, & u \sim v \\ 0, & \text{otherwise.} \end{cases}$$

Note that the Markov chain $\{X_n\}_{n \geq 0}$ is irreducible (that is, $\sum_{n=1}^{\infty} \mathbb{P}(X_n = v | X_0 = u) > 0$ for all $u, v \in V(G)$) and it is reversible (that is, there exists a nontrivial measure $\pi(dy)$ on $V(G)$, such that $\pi(u)q_{uv} = \pi(v)q_{vu}$ for all $u, v \in V(G)$). Indeed, irreducibility easily follows from connectedness of the graph G and for the reversibility measure we can take $\pi := c$. Also, let us remark that to every irreducible and reversible time-homogeneous Markov chain on a discrete state space S given by the transition kernel q_{uv} , $u, v \in S$, and a reversibility measure $\pi(dy)$ we can join a network $N = (G, c)$. Indeed, put $V(G) = S$, the vertices u and v are adjacent if $q_{uv} > 0$, the graph G is connected because of irreducibility of the corresponding Markov chain and the conductance is defined by $c(e_{uv}) = \pi(u)q_{uv}$.

Further, let $u_0 \in V(G)$ be an arbitrary vertex of the network N . A *flow* from u_0 to ∞ is a function $\theta : V(G) \times V(G) \rightarrow \mathbb{R}$ such that $\theta(u, v) = 0$ unless $u \sim v$, $\theta(u, v) = -\theta(v, u)$ for all

$u, v \in V(G)$ and $\sum_{v \in V(G)} \theta(u, v) = 0$ if $u \neq u_0$. We call the flow a *unit flow* if $\sum_{u \in V(G)} \theta(u_0, u) = 1$.

The *energy* of the flow is defined by

$$\mathcal{E}(\theta) = \frac{1}{2} \sum_{u \sim v} \frac{\theta^2(u, v)}{c(e_{uv})}.$$

Next, recall that a state u of a time-homogeneous Markov chain $\{X_n\}_{n \geq 0}$ on a discrete state space S is called *transient* if $\sum_{n=1}^{\infty} \mathbb{P}(X_n = u | X_0 = u) < \infty$ and it is called *recurrent* if $\sum_{n=1}^{\infty} \mathbb{P}(X_n = u | X_0 = u) = \infty$. If every state is transient (resp. recurrent) the chain itself is called transient (resp. recurrent). It is well known that every irreducible Markov chain is either transient or recurrent (see [MT93, Theorem 8.1.2]). Finally, the main tool for proving Theorem 2.1 is given in the following theorem.

Theorem 2.2. [Lyo83, Theorem 1] *Random walk on a network N is transient if and only if there is a unit flow on N of finite energy from some vertex to ∞ .*

Proof of Theorem 2.1. First, note that, according to [She62, Lemma 1.2], without loss of generality we can assume that $p_0 > 0$. Thus, $\{S_n\}_{n \geq 0}$ is a random walk on the network $N = (G, c)$, where $G = (V(G), E(G)) = (\mathbb{Z}, \mathbb{Z} \times \mathbb{Z})$ and $c(e_{uv}) = p_{|v-u|}$. Now, following the ideas from [HS02, Theorem 1], we construct a unit flow from 0 to ∞ for the random walk $\{S_n\}_{n \geq 0}$ such that the corresponding energy is bounded from above by (2.1). Then the desired result follows from Theorem 2.2. First, let us partition the set of vertices $V(G) = \mathbb{Z}$ on the sets $B_0 = \{0\}$, $B_i = \{2^{i-1}, 2^{i-1} + 1, \dots, 2^i - 1\}$ and $B_{-i} = \{-2^i + 1, \dots, -2^{i-1} - 1, -2^{i-1}\}$, $i \geq 1$, and let us define a unit flow $\theta : V(G) \times V(G) \rightarrow \mathbb{R}$ from 0 to ∞ in the following way. For $u \in B_i$ and $v \in B_j$ define

$$\theta(u, v) := \begin{cases} \frac{1}{2}, & i = 0 \text{ and } j = -1 \text{ or } j = 1 \\ 0, & i = j \text{ or } |i - j| \geq 2 \\ 2^{-2|i|}, & 0 < i < j = i + 1 \text{ or } j < j + 1 = i < 0. \end{cases}$$

Recall that flow has to be antisymmetric, hence we define $\theta(v, u) := -\theta(u, v)$. Next, note that

$$\sum_{v \in \mathbb{Z}} \theta(1, v) = \theta(1, 0) + \theta(1, 2) + \theta(1, 3) = -\frac{1}{2} + \frac{1}{4} + \frac{1}{4} = 0,$$

and analogously $\sum_{v \in \mathbb{Z}} \theta(-1, v) = 0$. Further, for $u \in B_i$, $i \geq 2$, we have

$$\sum_{v \in \mathbb{Z}} \theta(u, v) = \sum_{v \in B_{i-1}} \theta(u, v) + \sum_{v \in B_{i+1}} \theta(u, v) = -2^{i-2} 2^{-2(i-1)} + 2^i 2^{-2i} = 0,$$

and analogously for $u \in B_i$, $i \leq -2$, we have

$$\sum_{v \in \mathbb{Z}} \theta(u, v) = 0.$$

According to this, θ is a flow from 0 to ∞ . Finally, since $\sum_{v \in \mathbb{Z}} \theta(0, v) = \theta(0, 1) + \theta(0, -1) = 1$, θ is a unit flow from 0 to ∞ . Now, let us prove that the energy of the flow θ is bounded from above by

(2.1). We have

$$\begin{aligned}
\mathcal{E}(\theta) &= \frac{1}{2} \sum_{u \sim v} \frac{\theta^2(u, v)}{c(e_{uv})} = \frac{1}{2} \sum_{(u, v) \in E(G)} \frac{\theta^2(u, v)}{p_{|v-u|}} \\
&= \frac{1}{2} \sum_{u \geq 0, v \geq 0, u \neq v} \frac{\theta^2(u, v)}{p_{|v-u|}} + \frac{1}{2} \sum_{u \geq 0, v < 0} \frac{\theta^2(u, v)}{p_{|v-u|}} \\
&\quad + \frac{1}{2} \sum_{u < 0, v \geq 0} \frac{\theta^2(u, v)}{p_{|v-u|}} + \frac{1}{2} \sum_{u < 0, v < 0, u \neq v} \frac{\theta^2(u, v)}{p_{|v-u|}}.
\end{aligned}$$

Note that, from the symmetry of the distribution $\{p_n\}_{n \in \mathbb{Z}}$ and the definition of the flow θ , the second and the third term equal $\frac{1}{8p_1}$. Next, again from the symmetry of the distribution $\{p_n\}_{n \in \mathbb{Z}}$ and the symmetry of the function θ^2 , we have

$$\begin{aligned}
\mathcal{E}(\theta) &\leq \frac{1}{4p_1} + \sum_{u \geq 0, v \geq 0} \frac{\theta^2(u, v)}{p_{|v-u|}} \\
&= \frac{1}{4p_1} + \sum_{u \geq 0, w \geq 0} \frac{\theta^2(u, u+w)}{p_w} + \sum_{u \leq 0, w \leq 0} \frac{\theta^2(u, u+w)}{p_{|w|}} \\
&= \frac{1}{4p_1} + 2 \sum_{u \geq 0, w \geq 0} \frac{\theta^2(u, u+w)}{p_w}.
\end{aligned}$$

Note that $\theta(u, u+w) = 0$ when $u+w \geq 4u$, except for $u = 0$ and $w = 1$. Thus

$$\begin{aligned}
\mathcal{E}(\theta) &\leq \frac{3}{4p_1} + 2 \sum_{w \geq 2, u \geq \lceil \frac{w}{3} \rceil} \frac{\theta^2(u, u+w)}{p_w} \\
&= \frac{3}{4p_1} + \frac{1}{8p_2} + 2 \sum_{w \geq 2, u \geq \lceil \frac{w}{3} \rceil, u \neq 1} \frac{\theta^2(u, u+w)}{p_w},
\end{aligned}$$

where $\lceil x \rceil$ denotes the smallest integer not less than x . Now, since for $u \in B_i$, $i \geq 2$, (that is, for $u \geq 2$), we have $\theta^2(u, v) \leq (2^{-2(i-1)})^2 = 16(2^i)^{-4} \leq 16u^{-4}$, then

$$\sum_{u \geq \lceil \frac{w}{3} \rceil, u \neq 1}^{\infty} \theta^2(u, u+w) \leq \begin{cases} \int_{\lceil \frac{w}{3} \rceil - 1}^{\infty} \frac{16}{x^4} dx = \frac{16}{3(\lceil \frac{w}{3} \rceil - 1)^3} \leq \frac{16}{3(\frac{w}{3} - 1)^3} = \frac{144}{(w-3)^3}, & w \geq 4 \\ \int_1^{\infty} \frac{16}{x^4} dx = \frac{16}{3}, & w = 2, 3. \end{cases}$$

This yields

$$\begin{aligned}
\mathcal{E}(\theta) &\leq \frac{3}{4p_1} + \frac{1}{8p_2} + 2 \sum_{w \geq 2} \frac{1}{p_w} \sum_{u \geq \lceil \frac{w}{3} \rceil, u \neq 1} \theta^2(u, u+w) \\
&\leq \frac{3}{4p_1} + \frac{1}{8p_2} + \frac{32}{3p_2} + \frac{32}{3p_3} + 288 \sum_{w \geq 4} \frac{1}{(w-3)^3 p_w}.
\end{aligned}$$

This accomplishes the proof of Theorem 2.1. □

3 Continuous case

In this section, we prove Theorem 1.1 in the case of continuous jumps. As in the case of discrete jumps, the main step is to consider the random walk case.

Theorem 3.1. *Let $\{S_n\}_{n \geq 0}$ be one-dimensional symmetric random walk with jumps $\mathbb{P}(J_1 \in dy) = f(y)dy$, where the probability density function $f(y)$ is such that $f(y) > 0$ a.e. Then the random walk $\{S_n\}_{n \geq 0}$ is transient if*

$$\int_1^\infty \frac{dy}{y^3 f(y)} < \infty. \quad (3.1)$$

Again, similarly as in the case of discrete jumps, the transience condition for the Lévy process case can be easily derived from the random walk case. Indeed, let $\{L_t\}_{t \geq 0}$ be a one-dimensional symmetric Lévy process with the Lévy measure $\nu(dy) = f(y)dy$, where the density $f(y)$ is such that $f(y) > 0$ a.e. Then, first note that, according to [Sat99, Theorem 38.2], without loss of generality we can assume that $\nu(\mathbb{R}) < \infty$. Thus,

$$\{L_t\}_{t \geq 0} \stackrel{d}{=} \{S_{P_t}\}_{t \geq 0},$$

where $\{S_n\}_{n \geq 0}$ is a random walk with continuous jumps $\mathbb{P}(J_1 \in dy) := \frac{1}{\nu(\mathbb{R})}f(y)dy$ and $\{P_t\}_{t \geq 0}$ is the Poisson process with parameter $\nu(\mathbb{R})$ independent of $\{S_n\}_{n \geq 0}$. Now, the desired result follows from the definition of the transience in terms of sojourn times.

Before the proof of Theorem 3.1, we need some auxiliary results. Let $B \subseteq \mathbb{R}^d$ be an arbitrary Borel set and let us denote by $\mathbb{D}(\mathbb{R}^d)$ the space of \mathbb{R}^d -valued càdlàg functions equipped with the Skorohod topology. Define the *set of recurrent paths* by

$$R(B) := \{\omega \in \mathbb{D}(\mathbb{R}^d) : \forall n \in \mathbb{N}, \exists t \geq n \text{ such that } \omega(t) \in B\},$$

and the *set of transient paths* by

$$T(B) := \{\omega \in \mathbb{D}(\mathbb{R}^d) : \exists s \geq 0 \text{ such that } \omega(t) \notin B, \forall t \geq s\}.$$

In the following proposition, we characterize the transience and recurrence property of Lévy process in terms of càdlàg paths.

Proposition 3.2. *Let $\mathbf{L} = \{L_t\}_{t \geq 0}$ be an \mathbb{R}^d -valued Lévy process. Then, \mathbf{L} is transient if and only if $\mathbb{P}_{\mathbf{L}}(T(B_a)) = 1$ for all $a > 0$, and it is recurrent if and only if $\mathbb{P}_{\mathbf{L}}(R(B_a)) = 1$ for all $a > 0$, where B_a denotes the open ball of radius a around the origin.*

Proof. The proof follows directly from the definition of the transience and recurrence properties. \square

Now, let us recall the notion of characteristics of a semimartingale (see [JS03]). Let $(\Omega, \mathcal{F}, \{\mathcal{F}_t\}_{t \geq 0}, \mathbb{P}, \{S_t\}_{t \geq 0})$, $\{S_t\}_{t \geq 0}$ in the sequel, be a one-dimensional semimartingale and let $h : \mathbb{R} \rightarrow \mathbb{R}$ be a truncation function (that is, a continuous bounded function such that $h(x) = x$ in a neighborhood of the origin). We define two processes

$$\check{S}(h)_t := \sum_{s \leq t} (\Delta S_s - h(\Delta S_s)) \quad \text{and} \quad S(h)_t := S_t - \check{S}(h)_t,$$

where the process $\{\Delta S_t\}_{t \geq 0}$ is defined by $\Delta S_t := S_t - S_{t-}$ and $\Delta S_0 := S_0$. The process $\{S(h)_t\}_{t \geq 0}$ is a special semimartingale. Hence, it admits the unique decomposition

$$S(h)_t = S_0 + M(h)_t + B(h)_t, \quad (3.2)$$

where $\{S(h)_t\}_{t \geq 0}$ is a local martingale and $\{B(h)_t\}_{t \geq 0}$ is a predictable process of bounded variation.

Definition 3.3. Let $\{S_t\}_{t \geq 0}$ be a semimartingale and let $h : \mathbb{R} \rightarrow \mathbb{R}$ be the truncation function. Furthermore, let $\{B(h)_t\}_{t \geq 0}$ be the predictable process of bounded variation appearing in (3.2), let $N(\omega, ds, dy)$ be the compensator of the jump measure

$$\mu(\omega, ds, dy) = \sum_{s: \Delta S_s(\omega) \neq 0} \delta_{(s, \Delta S_s(\omega))}(ds, dy)$$

of the process $\{S_t\}_{t \geq 0}$ and let $\{C_t\}_{t \geq 0}$ be the quadratic co-variation process for $\{S_t^c\}_{t \geq 0}$ (continuous martingale part of $\{S_t\}_{t \geq 0}$), that is,

$$C_t = \langle S_t^c, S_t^c \rangle.$$

Then (B, C, N) is called the characteristics of the semimartingale $\{S_t\}_{t \geq 0}$ (relative to $h(x)$). If we put $\tilde{C}(h)_t := \langle M(h)_t, M(h)_t \rangle$, where $\{M(h)_t\}_{t \geq 0}$ is the local martingale appearing in (3.2), then (B, \tilde{C}, N) is called the modified characteristics of the semimartingale $\{S_t\}_{t \geq 0}$ (relative to $h(x)$).

Proposition 3.4. Let $\mathbf{S} = \{S_n\}_{n \geq 0}$ be a one-dimensional random walk with continuous jumps $\mathbb{P}(J_1 \in dy) = f(y)dy$. For $\delta > 0$, let $\mathbf{S}^\delta = \{S_n^\delta\}_{n \geq 0}$ be a random walk on $\delta\mathbb{Z}$ with discrete jumps

$$\mathbb{P}(J_1^\delta = \delta n) = \int_{\delta n - \frac{\delta}{2}}^{\delta n + \frac{\delta}{2}} f(y)dy, \quad n \in \mathbb{Z}.$$

Further, let $\{P_t\}_{t \geq 0}$ be the Poisson process with parameter 1 independent of \mathbf{S} and \mathbf{S}^δ , $\delta > 0$, and let $\bar{\mathbf{S}} := \{S_{P_t}\}_{t \geq 0}$ and $\bar{\mathbf{S}}^\delta := \{S_{P_t}^\delta\}_{t \geq 0}$. Then $\bar{\mathbf{S}}^\delta \xrightarrow{d} \bar{\mathbf{S}}$ when $\delta \rightarrow 0$, where \xrightarrow{d} denotes the convergence in $\mathbb{D}(\mathbb{R}^d)$ with respect to the Skorohod topology, and all the random walks \mathbf{S}^δ , $\delta > 0$, are either transient or recurrent at the same time and this transience and recurrence dichotomy is equivalent with the transience and recurrence dichotomy of the random walk \mathbf{S} .

Proof. Clearly, $\bar{\mathbf{S}}$ and $\bar{\mathbf{S}}^\delta$, $\delta > 0$, are processes of bounded variation. Thus, they are semimartingales. Further, let $h(x)$ be the truncation function and let (B, \tilde{C}, N) and $(B^\delta, \tilde{C}^\delta, N^\delta)$, $\delta > 0$, be the modified characteristics of $\bar{\mathbf{S}}$ and $\bar{\mathbf{S}}^\delta$, $\delta > 0$, respectively. Now, since $\bar{\mathbf{S}}$ and $\bar{\mathbf{S}}^\delta$ are Lévy processes, by [JS03, Proposition 2.17 and Corollary II.4.19], their (modified) characteristics are exactly the corresponding Lévy triplets, that is,

$$B_t = t\mathbb{E}[h(J_1)], \quad N(ds, dy) = ds\mathbb{P}(J_1 \in dy), \quad \tilde{C}_t = t\mathbb{E}[h^2(J_1)]$$

and

$$B_t^\delta = t\mathbb{E}[h(J_1^\delta)], \quad N^\delta(ds, dy) = ds\mathbb{P}(J_1^\delta \in dy), \quad \tilde{C}_t^\delta = t\mathbb{E}[h^2(J_1^\delta)].$$

According to this, in order to prove the desired convergence, by [JS03, Theorem VIII.2.17], it suffices to show that

$$\sup_{s \leq t} |B_s^\delta - B_s| \rightarrow 0, \quad \tilde{C}_t^\delta \rightarrow \tilde{C}_t \quad \text{and} \quad \int_{[0, t] \times \mathbb{R}} g(y)N^\delta(ds, dy) \rightarrow \int_{[0, t] \times \mathbb{R}} g(y)N(ds, dy)$$

when $\delta \rightarrow 0$ for all $t \geq 0$ and for every bounded and continuous function $g(x)$ vanishing in a neighborhood of the origin. Clearly, in order to prove the above convergences, it suffices to show that

$$\mathbb{E}[g(J_1^\delta)] \rightarrow \mathbb{E}[g(J_1)]$$

when $\delta \rightarrow 0$ for every bounded and continuous function $g(x)$. But this fact easily follows from [Bil99, Theorem 2.1], definition of the jumps $\{J_n^\delta\}_{n \geq 0}$, $\delta > 0$, and continuity of the jumps $\{J_n\}_{n \geq 0}$.

Now, we prove the second part of the proposition. Let $\delta_0 > 0$ be arbitrary. By completely the same arguments as above, we have $\bar{\mathbf{S}}^\delta \xrightarrow{d} \bar{\mathbf{S}}^{\delta_0}$ when $\delta \rightarrow \delta_0$. Next, let $a > 0$ be arbitrary, then

$T(B_a) = R(B_a)^c$, $R(B_a)$ is open in $\mathbb{D}(\mathbb{R})$ and $\partial R(B_a) \subseteq R(B_{a+\varepsilon}) \setminus R(B_{a-\varepsilon})$, where $\partial R(B_a)$ denotes the boundary of the set $R(B_a)$ and $0 < \varepsilon < a$. Thus, by Proposition 3.2, we have

$$\mathbb{P}_{\bar{\mathbf{S}}^{\delta_0}}(\partial R(B_a)) \leq \mathbb{P}_{\bar{\mathbf{S}}^{\delta_0}}(R(B_{a+\varepsilon})) - \mathbb{P}_{\bar{\mathbf{S}}^{\delta_0}}(R(B_{a-\varepsilon})) = 0$$

for all $a > 0$. Hence, the sets $T(B_a)$ and $R(B_a)$, $a > 0$, are continuity sets for $\mathbb{P}_{\bar{\mathbf{S}}^{\delta_0}}$. Now, by [Bil99, Theorem 2.1], this yields

$$\lim_{\delta \rightarrow \delta_0} \mathbb{P}_{\bar{\mathbf{S}}^\delta}(T(B_a)) = \mathbb{P}_{\bar{\mathbf{S}}^{\delta_0}}(T(B_a))$$

for all $a > 0$. Hence, for all $a > 0$, the function

$$\delta \mapsto \mathbb{P}_{\bar{\mathbf{S}}^\delta}(T(B_a))$$

is continuous on $(0, \infty)$. According to this, by Proposition 3.2, $\mathbb{P}_{\bar{\mathbf{S}}^\delta}(T(B_a)) = 1$ for all $\delta > 0$ and all $a > 0$, or $\mathbb{P}_{\bar{\mathbf{S}}^\delta}(T(B_a)) = 0$ for all $\delta > 0$ and all $a > 0$. This means, again by Proposition 3.2, that all the random walks $\bar{\mathbf{S}}^\delta$, $\delta > 0$, are either transient or recurrent at the same time.

Finally, by completely the same arguments as above, $\mathbb{P}_{\bar{\mathbf{S}}^\delta}(\partial T(B_a)) = 0$ for all $a > 0$. Then, again by [Bil99, Theorem 2.1], we have

$$\lim_{\delta \rightarrow 0} \mathbb{P}_{\bar{\mathbf{S}}^\delta}(T(B_a)) = \mathbb{P}_{\bar{\mathbf{S}}}(T(B_a))$$

for all $a > 0$. Thus, by Proposition 3.2, the transience and recurrence property of the random walks $\bar{\mathbf{S}}^\delta$, $\delta > 0$, is equivalent with the transience and recurrence property of the random walk $\bar{\mathbf{S}}$. \square

At the end, we prove Theorem 3.1.

Proof of Theorem 3.1. Let $\{S_n^1\}_{n \geq 0}$ be a random walk on \mathbb{Z} with discrete jumps

$$\mathbb{P}(J_1^1 = n) := \int_{n-\frac{1}{2}}^{n+\frac{1}{2}} f(y) dy, \quad n \in \mathbb{Z}.$$

Next, by the Jensen's inequality, we have

$$\int_{\frac{1}{2}}^{\infty} \frac{dy}{y^3 f(y)} = \sum_{n=1}^{\infty} \int_{n-\frac{1}{2}}^{n+\frac{1}{2}} \frac{dy}{y^3 f(y)} \geq \sum_{n=1}^{\infty} \frac{1}{(n+\frac{1}{2})^3 \int_{n-\frac{1}{2}}^{n+\frac{1}{2}} f(y) dy} = \sum_{n=1}^{\infty} \frac{1}{(n+\frac{1}{2})^3 \mathbb{P}(J_1^1 = n)}.$$

Thus, by Theorem 2.1, the random walk $\{S_n^1\}_{n \geq 0}$ is transient. Now, the desired result follows from Proposition 3.4. \square

4 Some remarks on the main results

In this section, we discuss some properties of the condition (1.1) we mentioned in Section 1. First, we prove that, under the assumption of unimodality, the condition (1.2) is stronger than the condition (1.1). Recall that a one-dimensional symmetric Lévy measure $\nu(dy)$ is unimodal if it is of the form $\nu(dy) = f(y)dy$, where the density function $f(y)$ is symmetric, decreasing on $(0, \infty)$ and it satisfies $\int_{|y|>\varepsilon} f(y)dy < \infty$ for all $\varepsilon > 0$. Let us fix $y_0 > 1$, then, by the Fubini's theorem, for all $y \geq y_0$ we

have

$$\begin{aligned}
\int_0^y z\nu(\max\{1, z\}, \infty) dz &= \int_0^y z \int_{\max\{1, z\}}^{\infty} f(u) du dz \\
&= \int_0^1 z \int_1^{\infty} f(u) du dz + \int_1^y z \int_z^{\infty} f(u) du dz \\
&= \frac{1}{2} \int_1^{\infty} f(u) du + \int_1^y z \int_z^{\infty} f(u) du dz \\
&= \frac{1}{2} \int_1^y u^2 f(u) du + \frac{y^2}{2} \int_y^{\infty} f(u) du \\
&\geq \frac{y^3 - 1}{6} f(y) + \frac{y^2}{2} \int_y^{\infty} f(u) du \\
&\geq Cy^3 f(y),
\end{aligned}$$

where in the fifth line we used the fact that $f(y)$ is decreasing on $(0, \infty)$ and $0 < C < \frac{y_0^3 - 1}{6y_0^3}$ is arbitrary. Now, we have

$$\begin{aligned}
&\int_1^{\infty} \left(\int_0^y z\nu(\max\{1, z\}, \infty) dz \right)^{-1} dy \\
&= \int_1^{y_0} \left(\int_0^y z\nu(\max\{1, z\}, \infty) dz \right)^{-1} dy + \int_{y_0}^{\infty} \left(\int_0^y z\nu(\max\{1, z\}, \infty) dz \right)^{-1} dy \\
&\leq \frac{2(y_0 - 1)}{y_0 \int_{y_0}^{\infty} f(y) dy} + \frac{1}{C} \int_{y_0}^{\infty} \frac{dy}{y^3 f(y)} \\
&\leq D \int_1^{\infty} \frac{dy}{y^3 f(y)},
\end{aligned}$$

for some suitably chosen constant $D > 0$. Therefore, we have proved the desired result.

Finally, we prove the transience property of a Lévy process with “multiple indices of stability” $\{L_t\}_{t \geq 0}$. Recall that $\{L_t\}_{t \geq 0}$ is a one-dimensional symmetric Lévy process with the Lévy measure $\nu(\{n\}) = p_n$, where $p_{2n} = (2n)^{-\alpha-1}$ and $p_{2n-1} = (2n-1)^{-\beta-1}$ for $n \geq 1$ and $\alpha, \beta \in (0, \infty)$. We claim that if $\alpha < 1$ and $\beta \geq 1$, then $\{L_t\}_{t \geq 0}$ is transient. Clearly, it suffices to consider the random walk case. Let $\{S_n\}_{n \geq 0}$ be a random walk on \mathbb{Z} with jumps $\mathbb{P}(J_1 = n) = c^{-1}p_n$, where $c := \sum_{n \in \mathbb{Z}} p_n$ is the norming constant and $p_n, n \geq 1$, are as above. First, let us define a sequence of stopping times $\{T_n\}_{n \geq 0}$ inductively by $T_0 := 0$ and

$$T_n := \inf\{k > T_{n-1} : S_k \in 2\mathbb{Z}\},$$

for $n \geq 1$, and let us prove that $\mathbb{P}(T_n < \infty) = 1$ for all $n \geq 1$. We have

$$\begin{aligned}
\mathbb{P}(T_1 = \infty) &= \mathbb{P}(S_k \in 2\mathbb{Z} + 1 \text{ for all } k \geq 1) \\
&= \lim_{k \rightarrow \infty} \mathbb{P}(S_l \in 2\mathbb{Z} + 1 \text{ for all } 1 \leq l \leq k) \\
&= \mathbb{P}(J_1 \in 2\mathbb{Z} + 1) \lim_{k \rightarrow \infty} (\mathbb{P}(J_1 \in 2\mathbb{Z}))^{k-1} = 0.
\end{aligned}$$

Now, let us assume that $\mathbb{P}(T_{n-1} < \infty) = 1$ and prove that $\mathbb{P}(T_n < \infty) = 1, n \geq 2$. Denote by $N := T_{n-1}$. Then, by the strong Markov property, we have

$$\begin{aligned}
\mathbb{P}(T_n < \infty) &= \mathbb{E}[1_{\{T_n < \infty\}}] = \mathbb{E}[1_{\{T_1 < \infty\}} \circ \theta_N] = \mathbb{E}[\mathbb{E}[1_{\{T_1 < \infty\}} \circ \theta_N | \mathfrak{F}_N]] \\
&= \mathbb{E}[\mathbb{E}^{S_N}[1_{\{T_1 < \infty\}}]] = \sum_{i \in \mathbb{Z}} \mathbb{E}[1_{\{S_N = 2i\}}] = 1,
\end{aligned}$$

where θ_n , $n \geq 0$, are the shift operators on the canonical state space $\mathbb{Z}^{\{0,1,2,\dots\}}$ defined by $(\theta_n \omega)(m) := \omega(n+m)$, $m \geq 0$, and $\mathcal{F}_N := \{A \in \mathcal{F} : A \cap \{N = n\} \in \sigma\{S_1, \dots, S_n\} \text{ for all } n \geq 1\}$. Thus, the Markov chain $X_n := S_{T_n}$ is well defined. Clearly, $\{X_n\}_{n \geq 0}$ is irreducible on $2\mathbb{Z}$. Further, note that $\{X_n\}_{n \geq 0}$ and $\{S_n\}_{n \geq 0}$ are transient or recurrent at the same time. Indeed, let us define the following stopping times $\tau := \inf\{n \geq 1 : S_n = 0\}$ and $\tilde{\tau} = \inf\{n \geq 1 : X_n = 0\}$. We have

$$\mathbb{P}(\tilde{\tau} = \infty) = \mathbb{P}(X_n \neq 0 \text{ for all } n \geq 1) = \mathbb{P}(S_n \neq 0 \text{ for all } n \geq 1) = \mathbb{P}(\tau = \infty).$$

Now, the desired result follows from [MT93, Propositions 8.1.3 and 8.1.4]. According to this, it suffices to prove the transience property of $\{X_n\}_{n \geq 0}$. Further, note that $\{X_n\}_{n \geq 0}$ is actually a symmetric random walk on $2\mathbb{Z}$. Indeed, for $n \geq 0$ and $i, j \in \mathbb{Z}$ we have

$$\begin{aligned} & \mathbb{P}(X_{n+1} = 2j | X_n = 2i) \\ &= \mathbb{P}(S_1 = 2j | S_0 = 2i) + \sum_{i_1 \in \mathbb{Z}} \mathbb{P}(S_1 = 2i_1 + 1 | S_0 = 2i) \mathbb{P}(S_2 = 2j | S_1 = 2i_1 + 1) + \dots \\ &= \mathbb{P}(S_1 = 2j - 2i | S_0 = 0) + \sum_{i_1 \in \mathbb{Z}} \mathbb{P}(S_1 = 2i_1 - 2i + 1 | S_0 = 0) \mathbb{P}(S_2 = 2j - 2i | S_1 = 2i_1 - 2i + 1) + \dots \\ &= \mathbb{P}(X_{n+1} = 2j - 2i | X_n = 0) \\ &= \mathbb{P}(X_1 = 2j - 2i). \end{aligned}$$

Thus, $\{X_n\}_{n \geq 0}$ is spatially homogeneous. Next, for $n, k \geq 0$ and $i, j \in \mathbb{Z}$ we have

$$\begin{aligned} \mathbb{P}(X_{n+k} - X_n = 2i) &= \sum_{j \in \mathbb{Z}} \mathbb{P}(X_{n+k} = 2i + 2j, X_n = 2j) \\ &= \sum_{j \in \mathbb{Z}} \mathbb{P}(X_{n+k} = 2i + 2j | X_n = 2j) \mathbb{P}(X_n = 2j) \\ &= \mathbb{P}(X_k = 2i), \end{aligned}$$

and for $k \geq 1$, $n_1, \dots, n_k \geq 0$, $0 \leq n_1 \leq \dots \leq n_k$, and $i_1, \dots, i_{k-1} \in \mathbb{Z}$ we have

$$\begin{aligned} & \mathbb{P}(X_{n_k} - X_{n_{k-1}} = 2i_{k-1}, \dots, X_{n_2} - X_{n_1} = 2i_1) \\ &= \sum_{j \in \mathbb{Z}} \mathbb{P}(X_{n_k} = 2i_{k-1} + \dots + 2i_2 + 2j, \dots, X_{n_2} = 2i_1 + 2j, X_{n_1} = 2j) \\ &= \sum_{j \in \mathbb{Z}} \mathbb{P}(X_{n_k} = 2i_{k-1} + \dots + 2i_2 + 2j | X_{n_{k-1}} = 2i_{k-2} + \dots + 2i_2 + 2j) \cdots \\ & \quad \mathbb{P}(X_{n_2} = 2i_1 + 2j | X_{n_1} = 2j) \mathbb{P}(X_{n_1} = 2j) \\ &= \mathbb{P}(X_{n_k - n_{k-1}} = 2i_{k-1}) \cdots \mathbb{P}(X_{n_2 - n_1} = 2i_1) \\ &= \mathbb{P}(X_{n_k} - X_{n_{k-1}} = 2i_{k-1}) \cdots \mathbb{P}(X_{n_2} - X_{n_1} = 2i_1). \end{aligned}$$

Symmetry is trivially satisfied. Thus, the claim follows. Finally, let us show that the random walk $\{X_n\}_{n \geq 0}$ is transient. For $i \in \mathbb{Z} \setminus \{0\}$ we have

$$\begin{aligned} \mathbb{P}(X_1 = 2i) &= \mathbb{P}(S_1 = 2i) + \sum_{j \in \mathbb{Z}} \mathbb{P}(S_1 = 2j + 1) \mathbb{P}(S_2 = 2i | S_1 = 2j + 1) + \dots \\ &\geq \mathbb{P}(S_1 = 2i) = c^{-1} p_{2i} = c^{-1} |2i|^{-\alpha-1}. \end{aligned}$$

Now, since $\alpha < 1$, from the condition (1.1) we have

$$\sum_{n=1}^{\infty} \frac{1}{(2n)^3 \mathbb{P}(X_1 = 2n)} < \infty.$$

Therefore, we have proved the desired result.

References

- [Bil99] P. Billingsley. *Convergence of probability measures*. John Wiley & Sons Inc., New York, second edition, 1999.
- [Dur10] R. Durrett. *Probability: theory and examples*. Cambridge University Press, Cambridge, fourth edition, 2010.
- [HS02] J. P. Hobert and J. Schweinsberg. Conditions for recurrence and transience of a Markov chain on \mathbb{Z}^+ and estimation of a geometric success probability. *Ann. Statist.*, 30(4):1214–1223, 2002.
- [JS03] J. Jacod and A. N. Shiryaev. *Limit theorems for stochastic processes*. Springer-Verlag, Berlin, second edition, 2003.
- [Lyo83] T. Lyons. A simple criterion for transience of a reversible Markov chain. *Ann. Probab.*, 11(2):393–402, 1983.
- [MT93] S. P. Meyn and R. L. Tweedie. *Markov chains and stochastic stability*. Springer-Verlag London Ltd., London, 1993.
- [Sat99] K. Sato. *Lévy processes and infinitely divisible distributions*. Cambridge University Press, Cambridge, 1999.
- [She62] L. A. Shepp. Symmetric random walk. *Trans. Amer. Math. Soc.*, 104:144–153, 1962.
- [She64] L. A. Shepp. Recurrent random walks with arbitrarily large steps. *Bull. Amer. Math. Soc.*, 70:540–542, 1964.