



## Response of *Lemna minor* L. to short-term cobalt exposure: The effect on photosynthetic electron transport chain and induction of oxidative damage



Lidija Begović<sup>a,1</sup>, Selma Mlinarić<sup>a,1</sup>, Jasenka Antunović Dunić<sup>a</sup>, Zorana Katanić<sup>a</sup>, Zdenko Lončarić<sup>b</sup>, Hrvoje Lepeduš<sup>c</sup>, Vera Cesar<sup>a,\*</sup>

<sup>a</sup> Josip Juraj Strossmayer University of Osijek, Department of Biology, Ulica cara Hadrijana 8/A, HR-31000 Osijek, Croatia

<sup>b</sup> Faculty of Agriculture, Josip Juraj Strossmayer University of Osijek, Ulica kralja Petra Svačića 1d, HR-31000 Osijek, Croatia

<sup>c</sup> Faculty of Humanities and Social Sciences, Josip Juraj Strossmayer University of Osijek, Lorenza Jägerova 9, HR-31000 Osijek, Croatia

### ARTICLE INFO

#### Article history:

Received 9 April 2015

Received in revised form 3 March 2016

Accepted 8 March 2016

Available online 10 March 2016

#### Keywords:

Cobalt

*Lemna minor*

Photosystem II

Oxygen evolving complex

Lipid peroxidation

### ABSTRACT

The effect of two concentrations of cobalt ( $\text{Co}^{2+}$ ) on photosynthetic activity and antioxidative response in *Lemna minor* L. were assessed 24, 48 and 72 h after the start of the exposure. Higher concentration of cobalt (1 mM) induced growth inhibition while lower concentration (0.01 mM) increased photosynthetic pigments content. Analysis of chlorophyll *a* fluorescence transients revealed high sensitivity of photosystem II primary photochemistry to excess of  $\text{Co}^{2+}$  especially at the higher concentration where decreased electron transport beyond primary quinone acceptor  $\text{Q}_\text{A}^-$  and impaired function of oxygen evolving complex (OEC) was observed. Due to impairment of OEC, oxygen production was decreased at higher  $\text{Co}^{2+}$  concentration. Activity of superoxide dismutase was mainly inhibited while lipid peroxidation increased, at both concentrations, indicating that cobalt-induced oxidative damage after short exposure and moreover, susceptibility of the membranes in the cell to cobalt toxicity. Results obtained in this study suggest possible application of used parameters as tools in assessment of early damage caused by metals.

© 2016 Elsevier B.V. All rights reserved.

### 1. Introduction

Excess uptake of metals, regardless of plant requirements, results in toxic effects. According to the report from the 2012 (EBRC and ARCHE, 2012) emission of cobalt in Europe to surface water was 90895.9 kg Co/year. Cobalt pollution can originate from anthropogenic sources through wastewater from industry, manufacture, processing, use and disposal of cobalt-containing products thus increasing environmental concentration constantly. The mean cobalt concentrations in rivers are ranging from 0.1 to 1.1  $\mu\text{g L}^{-1}$  (0.0017 to 0.019  $\mu\text{mol L}^{-1}$ ) while samples collected from surface water and groundwater near mining in USA contained cobalt

concentrations from <1 to 625 000  $\mu\text{g L}^{-1}$  (<0.017 to 0.01  $\mu\text{mol L}^{-1}$ ) (ATSDR, 2004).

Cobalt, as one of the major micronutrients in plants, is involved in the growth and metabolism (Palit et al., 1994). Mechanism of uptake and distribution of  $\text{Co}^{2+}$  in plants is species-dependent and differently regulated (Bakkaus et al., 2005). Toxic effect of cobalt was investigated on *Lemna minor* L. (Ince et al., 1999; Sree et al., 2015) as well as other plant species such as barley, oilseed rape and tomato (Nagajyoti et al., 2010). Results showed that  $\text{Co}^{2+}$  influences growth and biomass, chlorophyll content, starch accumulation, decreases photosynthetic efficiency, water potential and transpiration rate. Cobalt can displace Fe, Mn, Zn and Cu from physiologically important binding sites and thus might decrease uptake and translocation of essential micronutrients (Chatterjee and Chatterjee, 2000) and disable photochemical activity of the metal-containing pigments (Drzewiecka-Matuszek et al., 2005).

The photosystem II (PSII) and its component, the oxygen evolving complex (OEC), are the most sensitive to stress factors and are frequently used for evaluation of stress response in plants (Plekhanov and Chemeris, 2003). Capability of plants for adequate response when exposed to metal toxicity can be assessed by

\* Corresponding author.

E-mail addresses:

[lbegovic@biologija.unios.hr](mailto:lbegovic@biologija.unios.hr) (L. Begović), [smlinaric@biologija.unios.hr](mailto:smlinaric@biologija.unios.hr) (S. Mlinarić), [jantunovic@biologija.unios.hr](mailto:jantunovic@biologija.unios.hr) (J. Antunović Dunić), [zkatanic@biologija.unios.hr](mailto:zkatanic@biologija.unios.hr) (Z. Katanić), [zdenko.loncaric@pfos.hr](mailto:zdenko.loncaric@pfos.hr) (Z. Lončarić), [hlepedus@yahoo.com](mailto:hlepedus@yahoo.com) (H. Lepeduš), [vcesarus@yahoo.com](mailto:vcesarus@yahoo.com), [vera.cesar@biologija.unios.hr](mailto:vera.cesar@biologija.unios.hr) (V. Cesar).

<sup>1</sup> These authors have equally contributed to this paper.

variable chlorophyll *a* fluorescence (Appenroth et al., 2001; Joshi and Mohanty, 2004). Mohanty et al. (1989) proposed mechanism of cobalt action on  $Q_B$  function leading to the impairment of PSII activity at the  $Q_B$  site. Recent reports showed that beside the possibility to impair PSII functionality and electron transport, metals can also influence OEC functionality (Belatik et al., 2013; Hasni et al., 2013).

As a response to metal toxicity, plants activate their defense mechanisms to cope with enhanced reactive oxygen species (ROS) production which enables them to prevent or diminish further damage. Plants have an antioxidant defense system which includes numerous enzymatic and non-enzymatic components. Superoxide dismutase (SOD) plays important role in antioxidative defense catalyzing dismutation of superoxide ( $O_2^-$ ) to hydrogen peroxide ( $H_2O_2$ ) and oxygen ( $O_2$ ), therefore eliminating negative impact of ROS on the cell components and tissues. Metals are known to induce damage to cell membranes by lipid peroxidation, changing their structure and function. Lipid peroxidation is considered a reflection of oxidative stress in organism (Prasad, 2004).

*Lemna minor* L. can be found in numerous aquatic habitats and therefore represents a good biological indicator for detection of metal pollution. It is also frequently used as a model plant for studying physiological response to metal toxicity (Naumann et al., 2007; Panda, 2008; Razinger et al., 2008; Vidaković-Cifrek et al., 2015).

Previous studies have suggested that cobalt induces damage to the components of the photosynthetic apparatus hence influencing overall photosynthetic performance of the plant as well as activates SOD and elevates lipid peroxidation levels as a result of oxidative stress.

The aim of the present study was to investigate the early targets of cobalt toxicity by exploring concentration-dependent effect and changes induced in photosynthetic apparatus in *L. minor* along with possible induction of the oxidative stress. We used chlorophyll *a* fluorescence and oxygen evolution measurement to assess photosynthetic performance, measured activity of superoxide dismutase (SOD) and level of thiobarbituric acid reactive substances (TBARS) as a marker of lipid peroxidation.

## 2. Materials and methods

### 2.1. Plant material and cobalt treatment

*Lemna minor* L. plants were originally collected in the Botanical Garden of the Department of Biology, Faculty of Science, University of Zagreb in 1996 and sterilized according to Krajnčić and Devidé (1980). Stock cultures were grown under axenic conditions on the modified Pirson–Seidel's nutrient solution containing 3.95 mmol L<sup>-1</sup> KNO<sub>3</sub>, 5.46 mmol L<sup>-1</sup> CaCl<sub>2</sub> × 2H<sub>2</sub>O, 1.47 mmol L<sup>-1</sup> KH<sub>2</sub>PO<sub>4</sub>, 1.21 mmol L<sup>-1</sup> MgSO<sub>4</sub> × 7H<sub>2</sub>O, 49 μmol L<sup>-1</sup> Na<sub>2</sub>EDTA × 2H<sub>2</sub>O, 20 μmol L<sup>-1</sup> Fe-citrate, 1.5 μmol L<sup>-1</sup> MnCl<sub>2</sub> × 4H<sub>2</sub>O, 8.1 μmol L<sup>-1</sup> H<sub>3</sub>BO<sub>3</sub>, 29.2 mmol L<sup>-1</sup> sucrose and 0.66 mmol L<sup>-1</sup> asparagine at pH 4.55 (Pirson and Seidel, 1950). Stock cultures were generous gift from professor Vidaković-Cifrek (Faculty of Science, University of Zagreb, Croatia) and are maintained since 2006 in our laboratory.

For cobalt toxicity assessment, 8–11 colonies were inoculated in Erlenmeyer flasks filled with 70 mL of nutrient solution supplemented with CoCl<sub>2</sub> in the final concentration of 0.01 mM and 1 mM. Control plants were grown on modified Pirson–Seidel's nutrient solution. Plants, both control and treated, were maintained in growth chamber at 16/8 light/dark regime under white fluorescent lamps (Osram, Germany) with ~60 μmol photons m<sup>-2</sup> s<sup>-1</sup> at 22 ± 2 °C (Babić et al., 2009). For experimental purposes, colonies were harvested 24, 48 and 72 h after the start of the treatment.

### 2.2. Determination of mineral concentrations by ICP-OES

All samples were washed thoroughly with deionized water and then dried to constant weight at 60 °C. The dried samples were ground into a fine powder using a heavy metal-free ultra-centrifugal mill. All plant samples for measuring mineral concentrations were digested with 10 mL of a 5:1 mixture of HNO<sub>3</sub> and H<sub>2</sub>O<sub>2</sub> at 180 °C for 60 min in microwave oven (CEM Mars 6). The concentrations of minerals in solutions of digested plant samples were determined by inductively-coupled plasma optical emission spectroscopy, ICP-OES (PerkinElmer Optima 2100 DV). The whole series of plant samples (consisted of four replicates) were analyzed with an internal pooled plasma control and with the reference material prepared in the same way as the other samples. All samples were prepared and analyzed in duplicate.

### 2.3. Growth rate

Plant growth, expressed as growth rate, was monitored 24, 48 and 72 h after inoculation by counting all visible fronds in control and treated plants. Growth rate (GR) was calculated by using formula  $GR = \ln N_{t_i} - \ln N_{t_0} / t_i - t_0$  where  $N_{t_0}$  represents frond number at the beginning of the experiment ( $t_0$ ) and  $N_{t_i}$  represents frond number 24, 48 and 72 h after the start of the exposure ( $t_i$ ). Doubling time of frond number ( $T_d$ ) was calculated for 72 h using the equation  $T_d = \ln 2 / GR$  (Vidaković-Cifrek et al., 2013). To evaluate the effect of 0.01 mM and 1 mM Co<sup>2+</sup> on growth, percent inhibition of growth rate (%I<sub>r</sub>) was calculated using equation  $\%I_r = (GR_C - GR_T) / GR_C \times 100$  where  $GR_C$  represents average values of growth rate in control and  $GR_T$  represents average values of growth rate in treated plants.

### 2.4. Determination of photosynthetic pigment content

Fronds and roots were collected and homogenized in absolute acetone followed by extraction for 15 min at +4 °C and centrifugation for 10 min at 3000 rpm. Procedure was repeated until the plant material was completely uncolored. The chlorophyll and carotenoid content was determined spectrophotometrically. Concentrations of total chlorophylls (Chl *a* + *b*), chlorophyll *a* to chlorophyll *b* ratio (Chl *a/b*) and carotenoids (Car) were calculated according to Lichtenthaler (1987).

### 2.5. Fast chlorophyll *a* fluorescence measurement

Fast chlorophyll *a* fluorescence transients of randomly selected control and treated *L. minor* fronds were recorded with Plant Efficiency Analyzer (Handy PEA Hansatech Instruments Ltd. Norfolk, UK). All measurements were performed on fully dark-adapted fronds at room temperature. After 30 min of dark adaptation, the fronds were exposed to a pulse of saturating red light (3200 μmol m<sup>-2</sup> s<sup>-1</sup>, peak at 650 nm) and recorded fluorescence transients were analyzed using JIP test. The JIP test represents translation of the original recorded data to biophysical parameters that quantify the stepwise energy flow through PSII (Strasser et al., 2000, 2004). In normal physiological conditions, fluorescence transient goes from initial O ( $F_0$ ) to maximal P ( $F_m$ ) fluorescence intensity and it includes two intermediate steps, J step at approximately 2 ms and I step at approximately 30 ms.

In order to compare recorded OJIP transients two types of normalization were used. Double normalization between O and P steps was used for calculation of relative variable fluorescence  $W_{OP} = (F_t - F_0) / (F_P - F_0)$  and it is a measure of reduced  $Q_A$  fraction. Normalization between the steps O (50 μs) and J (2 ms) revealed variable fluorescence  $W_{OJ} = (F_t - F_0) / (F_J - F_0)$  named K-band (300 μs). It was given as variable fluorescence and plotted with difference

kinetics  $\Delta W_{OJ} = W_{OJ} - (W_{OJ})_{ref}$  in the 50  $\mu\text{s}$  – 2 ms time range. The values measured for *L. minor* plants without  $\text{Co}^{2+}$  treatment 24 h, 48 h and 72 h after the start of the exposure (Control) were used as referent value  $(W_{OJ})_{ref}$  (Strasser et al., 2004; Yusuf et al., 2010). Recorded data were:  $F_0$  (fluorescence intensity at 50  $\mu\text{s}$ ),  $F_m$  (maximal fluorescence intensity),  $F_{300}$  (fluorescence intensity at 300  $\mu\text{s}$ ),  $F_1$  (fluorescence intensity at 2 ms),  $F_2$  (fluorescence intensity at 30 ms) and  $t_{max}$  (time needed to reach  $F_m$ ). JIP test parameters calculated from the recorded values were: the maximum quantum yield of PSII ( $TR_0/ABS$ ), the probability that trapped exciton moves an electron further than  $Q_A$  ( $ET_0/TR_0$ ), the probability that absorbed photon moves an electron further than  $Q_A^-$  ( $ET_0/ABS$ ), the absorption flux per active reaction center ( $ABS/RC$ ), the trapping flux per active RC ( $TR_0/RC$ ), the electron transport flux per active RC ( $ET_0/RC$ ), the dissipation flux per active RC ( $DI_0/RC$ ). The performance index ( $PI_{ABS}$ ) was calculated using the density of RC on chlorophyll *a* basis ( $RC/ABS$ ), the flux ratio of trapping per dissipation ( $TR_0/DI_0$ ) and the electron transport beyond  $Q_A^-$  ( $ET_0/(TR_0 - ET_0^-)$ ). Detailed formulae and definitions of the used JIP-test parameters, according to Strasser et al. (2004) are shown in Table S1, Supplementary information.

## 2.6. Oxygen evolution measurement

Oxygen evolution was measured using the liquid-phase Clark-type oxygen electrode (Hansatech, UK) at 60  $\mu\text{mol photons m}^{-2} \text{s}^{-1}$ . All measurements were performed at constant temperature (25 °C). *L. minor* plants were placed in a chamber filled with bicarbonate buffer described by Prasad et al. (2001).

## 2.7. Measurement of superoxide dismutase activity

Measurement of superoxide dismutase activity (SOD, EC 1.15.1.1) was carried out according to Giannopolitis and Ries (1977) with modifications described by Lepeduš et al. (2011). Fronds and roots were ground using mortar and pestle in liquid nitrogen and approximately 200 mg of powdered tissue was used. One unit of SOD activity was defined as the amount of enzyme that inhibits the NBT photoreduction by 50%. Protein content in extracts was determined according to Bradford (1976) using bovine serum albumin as a standard.

## 2.8. Measurement of lipid peroxidation level

Plant material, fronds and roots, was ground using mortar and pestle in liquid nitrogen and about two hundred milligrams of fine powder was used for analyses. The level of lipid peroxidation was determined as described by Verma and Dubey (2003) with modifications according to Lepeduš et al. (2011). The concentration of lipid peroxidation products were expressed as total thiobarbituric acid reactive substances level (TBARS) in nmol per g of fresh weight using an extinction coefficient of 155  $\text{mM}^{-1} \text{cm}^{-1}$ .

## 2.9. Statistical analysis

Statistical analyses were performed between control and treated plants using one-way analysis of variance (ANOVA) followed by post hoc LSD test for each exposure time separately, if not stated otherwise. Results were expressed as arithmetic means of 9 replicates from three experiments  $\pm$  standard deviation (SD) except for measurement of chlorophyll *a* fluorescence when 50 replicates were used and ICP-OES analysis when eight replicates were used. Different letters (a, b, c) represent statistical difference at  $P \leq 0.05$ . All statistical analyses were conducted with Statistica 12.5 Software (StatSoft Inc., USA).




Fig. 1. Growth rate of *L. minor* 24, 48 and 72 h after the start of the exposure to 0.01 mM and 1 mM  $\text{CoCl}_2$ . Analyses by ANOVA showed significant differences between control and treated plants within each exposure time. Values are represented as means  $\pm$  SD ( $n=9$ ).

## 3. Results and discussion

### 3.1. Cobalt content and influence on growth rate in *L. minor*

Results of cobalt content determination (Table 1) showed significant difference between control and treated plants and between two cobalt concentrations (0.01 and 1 mM). In vascular plants, growing in unpolluted freshwater, mean cobalt concentrations are 0.32 – 0.37  $\mu\text{g g}^{-1}$  dry weight (0.005–0.006  $\mu\text{mol g}^{-1}$  dry weight) (Outridge and Noller, 1991). These results are in concordance with measured  $\text{Co}^{2+}$  concentrations in control plants (Table 1). In plants treated with  $\text{CoCl}_2$  significant increase in accumulation, up to  $10^4$ -fold higher than in the control, approximately 3500  $\mu\text{g g}^{-1}$  dry weight (58.7  $\mu\text{mol g}^{-1}$  dry weight), can be observed 72 h after the start of the exposure to the higher concentration (1 mM) while at the lower concentration (0.01 mM) cobalt content was about 200-fold higher than in the control, 45.45  $\mu\text{g g}^{-1}$  dry weight (0.771  $\mu\text{mol g}^{-1}$  dry weight). In Sree et al. (2015) accumulation rate and consequent cobalt content was much higher. The authors reported approximately 900  $\mu\text{g g}^{-1}$  fresh weight of cobalt content after 72 h and 21  $\text{mg g}^{-1}$  dry weight was measured after 7 days in *L. minor* treated with the same  $\text{Co}^{2+}$  concentration as we used (0.01 mM).

Further, the authors reported on significant decrease in relative growth rate after seven days, up to 56% in plants treated with 0.01 mM and 93% in plants treated with 0.1 mM  $\text{Co}^{2+}$  (Sree et al., 2015). In our study the earliest influence of  $\text{Co}^{2+}$  on growth was observed 24 h after the start of the exposure, but only at higher concentration (1 mM) which caused inhibition of relative growth rate ( $\%I_r$ ) by 53% compared to control. This effect was also seen 72 h after the start of the exposure when significant inhibition (63%) of relative growth rate (0.102  $\text{d}^{-1}$ ) with doubling time of 7.14 was observed (Fig. 1). Lower cobalt concentration (0.01 mM) did not influence relative growth rate (0.279  $\text{d}^{-1}$ ) at any exposure time as compared to control (Fig. 1). Doubling time at the lower concentration for 72 h was 2.55. In control plants relative growth rate for 72 h was 0.277  $\text{d}^{-1}$  with a doubling time of 2.59.

Ince et al. (1999) showed that higher concentrations of  $\text{Co}^{2+}$  inhibited growth in *L. minor* while lower  $\text{Co}^{2+}$  concentration stimulated growth. Stimulating effect of cobalt on growth rate was also observed in algae (El-Sheekh et al., 2003; Horvatić and Peršić, 2007; Osman et al., 2004) and sweet pepper (Gad and Hassan, 2013). However, cobalt-induced stimulation of growth was not observed in our study.

**Table 1**  
Cobalt and manganese content ( $\mu\text{g g}^{-1}$  dry weight  $\pm$  SD) in *L. minor* plants 24, 48 and 72 h after the start of the exposure to 0.01 and 1 mM concentration of  $\text{CoCl}_2$ . Student's *t*-test was used to analyze statistical difference between control and treated plants ( $n = 8$ ).

	$\text{Mn}^{2+}$					
	24 h		48 h		72 h	
Control	0.15 $\pm$ 0.03 (0.003 $\pm$ 0.001)	0.25 $\pm$ 0.08 (0.004 $\pm$ 0.001)	0.38 $\pm$ 0.10 (0.007 $\pm$ 0.002)	87.33 $\pm$ 8.59 (1.592 $\pm$ 0.137)	91.90 $\pm$ 12.09 (1.676 $\pm$ 0.220)	94.23 $\pm$ 1.49 (1.718 $\pm$ 0.082)
0.01 mM	40.11 $\pm$ 3.41* (0.68 $\pm$ 0.058)	52.03 $\pm$ 1.97* (0.883 $\pm$ 0.033)	45.45 $\pm$ 7.63* (0.771 $\pm$ 0.129)	80.25 $\pm$ 3.25* (1.463 $\pm$ 0.059)	86.87 $\pm$ 11.97* (1.584 $\pm$ 0.218)	82.19 $\pm$ 7.75* (1.498 $\pm$ 0.141)
1 mM	2037.00 $\pm$ 11.31* (34.566 $\pm$ 3.908)	2646.30 $\pm$ 467.74* (44.905 $\pm$ 7.937)	3459.00 $\pm$ 202.78* (58.697 $\pm$ 3.441)	63.10 $\pm$ 6.69* (1.150 $\pm$ 0.122)	58.79 $\pm$ 11.70* (1.072 $\pm$ 0.213)	41.15 $\pm$ 3.16* (0.750 $\pm$ 0.112)

Asterisks indicate values that are significantly different from control plants ( $P \leq 0.05$ ). Values in parenthesis are expressed as ( $\mu\text{mol g}^{-1}$  dry weight  $\pm$  SD).

Differences, such as cobalt content and effect on growth could be due to different mechanisms that algae and different plant species use to absorb ions. In addition, different *L. minor* clones, culture medium and growth conditions used in previous studies (Ince et al., 1999; Sree et al., 2015) might have affected the accumulation rate of cobalt and caused difference in the effect of cobalt on growth. Also, prolonged exposure time must not be overlooked.


### 3.2. Influence of cobalt on photosynthetic pigments and mineral content

Morphological changes, like chlorosis, are usually associated with metal toxicity (Assche and Clijsters, 1990; Chatterjee and Chatterjee, 2003; Khan and Khan, 2010; Tkalec et al., 2008). As described in the paper by Sree et al. (2015) younger fronds of *L. minor*, as opposed to mother fronds, showed chlorotic or necrotic features after exposure to 0.01 mM of  $\text{Co}^{2+}$ , while at higher concentration (0.1 mM) mother fronds were more prone to the damage than younger ones since their growth was evidently inhibited. In our study, after exposure to 0.01 mM and 1 mM of  $\text{Co}^{2+}$ , plants did not show visible morphological changes indicating that longer exposure time is needed to induce such changes or, as mentioned before, it could be because of the different cobalt content measured in plants upon exposure to 0.01 mM of  $\text{Co}^{2+}$ . Previous research has shown that cobalt can inhibit or induce synthesis of photosynthetic pigments depending on the concentration. Significant increase in total chlorophyll and carotenoid content was observed 24, 48 and 72 h after the start of treatment with 0.01 mM concentration compared to control plants (Fig. 2A and B). Increase in the ratio of chlorophyll *a* and *b* was also observed but only 24 h after treatment with 0.01 mM concentration of  $\text{Co}^{2+}$  (Fig. 2C). In the study by Sree et al. (2015) chlorophyll and carotenoid content did not change after four days of exposure to 0.01 mM  $\text{Co}^{2+}$ . However, our results showed that lower (0.01 mM) concentration of  $\text{Co}^{2+}$  stimulated production of chlorophylls and carotenoids, which could be associated with the absence of chlorosis in treated plants.

Stimulation of chlorophyll production was also reported by Csartoday et al. (1984) in algae *Anacystis nidulans*, Zeid (2001) in bean and Tewari et al. (2002) in mung bean. Authors suggested that increase of chlorophyll content is a result of enhanced accumulation of protoporphyrin XI, one of the precursors in chlorophyll biosynthesis.

On the other hand, some earlier studies have reported decreased chlorophyll content, when higher concentrations of cobalt were applied, as a result of the inhibitory effect of cobalt on chlorophyll synthesis (Chatterjee and Chatterjee, 2000; Csartoday et al., 1984; Tewari et al., 2002). In our study, higher concentration of  $\text{Co}^{2+}$  did not decrease total chlorophyll and carotenoid content when compared to control plants (Fig. 2A and B). Therefore, to achieve inhibition of chlorophyll and carotenoid synthesis a longer period of exposure is required as demonstrated by Sree et al. (2015) whose treatment with 0.01 mM and 0.1 mM  $\text{Co}^{2+}$  induced a decrease after seven days.

Zn content was not affected by cobalt treatment (Table S2). Analyses of Fe content revealed differences between control and treated plants. Results of Sree et al. (2015) showed that Fe content did not change in *L. minor* plants upon exposure especially to 0.01 mM  $\text{Co}^{2+}$  after 1 and 7 days. Although in our study Fe content was not significantly different between control and 0.01  $\text{Co}^{2+}$  after 24 h, significant reduction of Fe content was observed 72 h after the start of the exposure to both concentrations of cobalt (Table S2). The main difference between our results and those of Sree et al. (2015) is due to the fact that in our control plants Fe content continuously increased for 72 h, while in the study by Sree et al. (2015) this trend was absent. As discussed previously, composition of nutrient medium,



**Fig. 2.** Total chlorophyll content, Chl a + b, (A), carotenoid content, Car (B), and chlorophyll a to b ratio, Chl a/b (C) in *L. minor* 24, 48 and 72 h after the start of the exposure to 0.01 mM and 1 mM concentrations of  $\text{Co}^{2+}$ . Significant differences between control and treated plants were designated by different letters (ANOVA,  $P \leq 0.05$ ). Values are represented as means  $\pm$  SD ( $n=9$ ).

light intensity as well as different clones could have influenced Fe content.

Mg content in *L. minor* was not affected by treatment with 0.01 mM  $\text{Co}^{2+}$  whereas the higher concentration caused a reduction already 24 h after the start of the exposure and the decrease was observed up to 72 h (Table S2). We can assume that over the time  $\text{Co}^{2+}$  would inhibit the chlorophyll formation by disabling incorporation of Mg, confirming the suggestion by Sree et al. (2015) and Shalygo et al. (1999)


### 3.3. Fast chlorophyll a fluorescence transients after short exposure to $\text{Co}^{2+}$

Detail analysis of fast chlorophyll a fluorescence transients showed changes in functioning of photosynthetic apparatus of *L. minor* plants exposed to cobalt. Difference could be observed between treated and untreated plants, as well as between plants treated with different concentrations of cobalt starting from 24 h onwards. Both control and treated plants had typical chlorophyll a fluorescence rise kinetics consisting of clearly visible O, J, I and P


steps indicating the ability for photosynthesis upon short exposure to cobalt.

In order to compare the effect of two concentrations of  $\text{Co}^{2+}$  we plotted the curves as a relative variable fluorescence derived from double O-P normalized transients ( $W_{OP} = (F_t - F_0)/(F_P - F_0)$ ). The results showed that 1 mM concentration increased J and I step in treated plants whereas 0.01 mM concentration decreased them when compared to control. The observed changes were statistically significant (Fig. 3A-C). Rise of  $V_j$  (Table S3, Supplementary information) suggests an increase of the proportion of closed PSII RCs and of the proportion of reduced  $Q_A$  at J step. The increase of  $V_i$  (Table S3, Supplementary information) indicates accumulation of the reduced  $Q_A$  and plastoquinone which cannot transfer electrons further than  $Q_A$  (Strasser et al., 2004). Similar results were obtained by Pan et al. (2010) in maize treated with antimony (Sb). Lower fluorescence level on J and I step was the result of lower accumulation rate of the reduced  $Q_A$  electron acceptor.

Both tested concentrations of cobalt had negative effect on PSII primary photochemistry measured as a decrease of  $\text{TR}_0/\text{ABS}$  (Fig. 4). However, while treatment with the higher cobalt concentration led



**Fig. 3.** Chlorophyll *a* fluorescence transients (OJIP) of dark adapted *L. minor* fronds in control plants and plants exposed to 0.01 mM and 1 mM  $\text{Co}^{2+}$  for 24 (A), 48 (B) and 72 h (C). Results are represented as double, O-P normalized transients ( $W_{OP} = (F_t - F_0) / (F_P - F_0)$ ). Native curves are shown as inserts in each panel. Each curve represents average kinetics of 50 replicates ( $n = 50$ ). The O, J, I and P steps each curve are marked in the plots.



**Fig. 4.** Photosynthetic parameters determined by the JIP-test analysis of fluorescence transients. Every plot consists of 13 structural and functional photosynthetic parameters (average value of 50 replicates) derived by the JIP-test from the fluorescence transients. Values were normalized using the control treatment (without symbol) as reference. Normalization was obtained for each parameter, 24 (A), 48 (B) and 72 h (C) after the start of  $\text{Co}^{2+}$  treatment. Statistical analysis of presented parameters is shown in Table S3.

to decreased overall photosynthetic efficiency expressed as  $PI_{ABS}$ , treatment with the lower cobalt concentration had the opposite effect inducing an increase of  $PI_{ABS}$  value (Fig. 4).

$PI_{ABS}$  is calculated as the product of three parameters:  $RC/ABS$ ,  $TR_0/DI_0$  and  $ET_0/(TR_0 - ET_0)$ . Changes of  $ET_0/(TR_0 - ET_0)$  were shown to be the main reason for the observed differences in  $PI_{ABS}$ . Comparison of several other parameters describing the efficiency of electron transport such as  $ET_0/TR_0$  or  $ET_0/ABS$  were decreased upon 1 mM cobalt treatment and increased upon 0.01 mM cobalt treatment. This indicated that differential effect of tested cobalt concentration on photosynthesis of *L. minor* is associated with effect on electron transport from reduced  $Q_A^-$  further downstream


(Fig. 4). Similar results were obtained by Perreault et al. (2014) who observed a biphasic response in electron transport in plants exposed to copper. This conclusion is further supported by analysis of specific energy fluxes describing the distribution of energy through PSII at the reaction center level. The lower concentration of  $\text{Co}^{2+}$  decreased the value which expresses the rate by which an excitation is trapped by the RC resulting in the reduction of  $Q_A$  to  $Q_A^-$  ratio ( $\text{TR}_0/\text{RC}$ ) but increased reoxidation of reduced  $Q_A^-$  via electron transport in an active RC ( $\text{ET}_0/\text{RC}$ ) at 24 and 48 h (Table S3, Supplementary information). Also, lower concentration of  $\text{Co}^{2+}$  negatively affected antenna size of active reaction centers 24 and 72 h after the start of the exposure causing a decrease in ABS/RC ratio (Table S3, Supplementary information). The ABS/RC value expresses the total absorption of PSII antenna chlorophylls divided by the number of active (in the sense of  $Q_A$  reducing) reaction centers. Thus, it is influenced by the ratio of active/inactive RCs (Falqueto et al., 2013). An increase in ABS/RC induced by high  $\text{Co}^{2+}$  concentration could be due to inactivation of RCs that are being transferred to non- $Q_A$ -reducing centers and/or due to increase in functional antenna size (Yusuf et al., 2010). Since the increase in ABS/RC in *L. minor* at 1 mM of  $\text{Co}^{2+}$  24 h after the start of the exposure was accompanied by a decrease in  $\text{TR}_0/\text{ABS}$  (Fig. 4), but  $\text{TR}_0/\text{RC}$  (Table S3, Supplementary information) was not affected, it can be assumed that some part of RCs was inactivated. Possible causes of RC inactivation are inactivation of OEC and/or structural alteration of heat sinks (or silent centers) (Yusuf et al., 2010). In our case, both events occurred. Heat sinks are able to dissipate excess of absorbed energy safely as heat instead of utilizing it to reduce  $Q_A$  and presence of those is indicated by the increase in  $\text{DI}_0/\text{RC}$  (Table S3, Supplementary information).

These results indicated that exposure of *L. minor* to 0.01 mM concentration could be limited to PSII, at least for the first 72 h of exposure while 1 mM concentration could have more extensive effect on the whole photosynthetic electron transport chain, from donor side of PSII up to the reduction of end acceptor of PSI. Results observed after treatment with higher concentration of cobalt are in agreement with those obtained by Mohanty et al. (1989).

### 3.4. Influence of cobalt on OEC

Appearance of positive K-band, which is considered to reflect inactivation of OEC (especially of the Mn complex) and/or increase the functional PSII antenna size (Yusuf et al., 2010), has been mainly used as an indicator of drought stress (Oukarroum et al., 2007) and heat stress in plants before visible symptoms appear (Lazár et al., 1997; Srivastava et al., 1997). Results of our investigation showed that K-band could be used as an indicator in the assessment of early stress caused by cobalt, particularly intactness of OEC. Although hidden in OJIP transients, further analyses using data normalized between O and J steps (K-band) expressed as variable fluorescence  $W_{OJ} = (F_t - F_0)/(F_J - F_0)$  and then plotted with difference kinetics  $\Delta W_{OJ} = W_{OJ} - (W_{OJ})_{\text{ref}}$  in the 50  $\mu\text{s}$ –2 ms time range. Arithmetic mean from values measured in control plants was used as referent value  $(W_{OJ})_{\text{ref}}$ . Lines (solid, dashed and dotted) refer to difference kinetics ( $\Delta W_{OJ}$ ) curves, while solid lines with marks (open and closed) refer to variable kinetics ( $W_{OJ}$ ) curves.


Reports of cobalt influence on plants shows that cobalt has the ability to displace metal ions from protein complexes as well as reducing their translocation from root to shoot (Chatterjee and Chatterjee, 2000). In studies on green microalgae *Scenedesmus* (Mallick and Mohn, 2003) and poplar (Velikova et al., 2011) exposed to metal stress authors suggested that the metals used replace Mn ion from OEC. Dissociation of a manganese-stabilizing protein, bound to the donor side of the reaction center of PS II and the release of Mn atoms is associated with the loss of manganese cluster (Brestic and Zivcak, 2013) and therefore OEC inactivation. Indeed, analyses of manganese content in treated plants revealed



**Fig. 5.** Changes in the shape of the chlorophyll *a* fluorescence transient curves of *L. minor* exposed to 0.01 mM and 1 mM  $\text{Co}^{2+}$  24 (A), 48 (B) and 72 h (C) after the start of exposure. Fluorescence data (average value of 50 replicates) are normalized between O and J steps (K-band) expressed as variable fluorescence  $W_{OJ} = (F_t - F_0)/(F_J - F_0)$  and then plotted with difference kinetics  $\Delta W_{OJ} = W_{OJ} - (W_{OJ})_{\text{ref}}$  in the 50  $\mu\text{s}$ –2 ms time range. Arithmetic mean from values measured in control plants was used as referent value  $(W_{OJ})_{\text{ref}}$ . Lines (solid, dashed and dotted) refer to difference kinetics ( $\Delta W_{OJ}$ ) curves, while solid lines with marks (open and closed) refer to variable kinetics ( $W_{OJ}$ ) curves.

significant decrease in comparison with control plants confirming the hypothesis of cobalt action on Mn ions (Table 1).

Described limitation on donor side of PSII under the influence of cobalt was not observed when lower cobalt concentration was applied in nutrient solution. On the contrary, occurrence of negative K-band was proved (Fig. 5A–C). Negative K-band has previously been demonstrated as an indicator of better performance of plants under stress conditions (De Ronde et al., 2004; Tewari et al., 2002; Yusuf et al., 2010). In this study it could indicate a mechanism by which the plant is trying to diminish the negative effect of small excess of cobalt caused by 0.01 mM  $\text{Co}^{2+}$  treatment.




**Fig. 6.** Influence of two cobalt concentrations (0.01 and 1 mM) on oxygen production in *L. minor* 24, 48 and 72 h after the start of exposure. Significant differences between control and treated plants were designated by different letters (ANOVA,  $P \leq 0.05$ ). Values are given as means  $\pm$  SD ( $n = 9$ ).

The conclusion about the above described influence of cobalt on OEC was further supported by oxygen evolution measurement (Fig. 6). The exposure to higher  $\text{Co}^{2+}$  concentration caused a decrease in oxygen production confirming inactivation of OEC indicated by the positive K-band (Fig. 5A–C), while lower  $\text{Co}^{2+}$  concentration increased oxygen production after 72 h (Fig. 6C).

### 3.5. SOD activity and TBARS levels in *L. minor* after exposure to $\text{Co}^{2+}$

Although SOD is the first line of antioxidative defense in plants and is therefore an important biomarker of oxidative stress, in our study SOD was not activated. On a contrary, SOD activity decreased in plants treated with 0.01 mM cobalt concentration 24, 48 and 72 h after the start of the exposure and 24 and 72 h after the start of the exposure to 1 mM concentration of cobalt (Fig. 7A). Increased activity of SOD was observed 48 h after the start of the exposure to the higher concentration of  $\text{Co}^{2+}$  (Fig. 7A). The higher value of SOD activity measured in control plants 24 h after the start of the exposure (Fig. 7A) could be caused by activation of antioxidative enzymes as a part of the unspecific reaction/response to changes

in the environment. Namely, according to Lichtenthaler (1996) in order to acclimate to the new conditions plants can activate their stress-coping mechanism which include different metabolic processes. This is so called general acclimatization syndrome (GAS) that includes initial oxidative burst, i.e. biologically unspecific reaction. On the other hand, prolonged and continuous stress will lead to damage or activation of defense mechanisms. Accordingly, in our study, the higher values of SOD activity, measured in control plants (24 h), could be a result of unspecific signaling as a response to inoculation to the nutrient medium prior to the beginning of the experiment. This statement is corroborated by the obtained results at 48 and 72 h which show lower SOD activities in control plants. It would be expected that treated plants also exhibit higher SOD activity after 24 h because of unspecific signaling caused by mechanical stress, but this was not the case in our study. Such result could be explained by the fact that increased *in vivo* cobalt content in *L. minor* was detectable as early as ten minutes after the start of the exposure demonstrating high accumulation rate (Sree et al., 2015). Under these conditions the amount of  $\text{Co}^{2+}$  ions present in the nutrient medium was most likely sufficient enough to reduce the SOD activity in treated plants.



**Fig. 7.** Specific activity of SOD (A) and TBARS content (B) in *L. minor* 24, 48 and 72 h after the start of the exposure to 0.01 mM and 1 mM  $\text{CoCl}_2$ . Significant differences between control and treated plants were designated by different letters (ANOVA,  $P \leq 0.05$ ). Values are represented as means  $\pm$  SD ( $n = 9$ ).



Decreased activity of SOD, in plants exposed to cadmium stress, was observed in pea (Sandalo et al., 2001) and wheat (Milone et al., 2003). In pigeon pea plants treated with cobalt Gopal (2014) reported a reduced SOD activity with increased concentration. The author suggested that decreased SOD activity is due to inhibition of SOD synthesis or inactivation of particular SOD isoforms as well as inactivation of H<sub>2</sub>O<sub>2</sub>. Because Co<sup>2+</sup> has affinity for complexes that possess Fe, Mn, Zn and Cu, it can switch place with these metals, whereby the activity of enzymes possessing them is reduced (Chatterjee and Chatterjee, 2000). In addition, inhibition of electron transport further than Q<sub>B</sub> could diminish production of superoxide anion (O<sub>2</sub><sup>-</sup>) production at PSI site (Chen et al., 2012; Pospíšil, 2012) which could be a consequence of decreased SOD activity in our study.

Action of ROS on the cell membrane results in the process of peroxidation of polyunsaturated fatty acids of lipids leading to lipid peroxidation (Wojtaszek, 1997). Previous studies indicated differential impact of Co<sup>2+</sup> on level of lipid peroxidation in treated plants. While Gopal (2014) reported on reduced lipid peroxidation level in pigeon pea, Zeid et al. (2013) reported increased TBARS content in alfalfa treated with increasing concentrations of cobalt. The latter is in agreement with our results. Exposure of *L. minor* to 1 mM of Co<sup>2+</sup> significantly increased lipid peroxidation levels (TBARS) already 24 h after the start of the exposure, compared to control (Fig. 7B). This increment was observed at 48 and 72 h also. Lipid peroxidation levels in plants treated with 0.01 mM concentration of Co<sup>2+</sup> significantly increased only 72 h after the start of the exposure, compared to control, but this increase was lower when compared to 1 mM concentration. Increased TBARS levels 24 h after the start of the exposure showed that higher concentration of cobalt induced early oxidative damage in *L. minor* while at lower concentration plant is able to prevent oxidative damage for some time, but the negative effect of cobalt is evident 72 h after the start of the treatment.

#### 4. Conclusion

It is important to identify how metals impact biochemical pathways in plants before visible toxic effects take place. Induction of defense mechanisms like antioxidative response and adjustment of photosynthetic performance enables the plant to protect itself when exposed to metals. In this study, toxic effects of two cobalt concentrations on *L. minor* photosynthetic activity and oxidative damage after short exposure were demonstrated. Both concentrations of cobalt mainly prevented the activation of SOD. On the other hand, TBARS content increased 24 h after the start of the exposure to 1 mM Co<sup>2+</sup> and 72 h after the start of the exposure to 0.01 mM indicating oxidative damage and showing that the integrity of the membranes in the cell was compromised. Therefore, the lipid peroxidation level could be a good marker for an early assessment of Co<sup>2+</sup> toxicity. Our results confirmed that Co<sup>2+</sup> influences photosynthetic performance altering the electron transport through PSII and induces OEC impairment, particularly at the higher concentration. At the lower concentration plants were increasing photosynthetic electron transport trying to adapt and to reduce the toxic effect of cobalt.

In toxicological studies it is important to detect physiological parameters of early stress in order to reduce or prevent the harmful consequences of metal toxicity. In the present study, chlorophyll a fluorescence parameters, especially K-band, were shown to be useful tools for assessment and screening of early physiological stress in *L. minor* induced by cobalt. Further investigations should be undertaken to validate the application of K-band for assessment of early toxic effect of other metals.

#### Acknowledgements

We would like to thank Dr. Vidaković-Cifrek, Faculty of Science, University of Zagreb, Croatia, for the valuable advice and suggestions. This study was supported by the Ministry of Science, Education and Sports, Republic of Croatia (grants no. 073-0731674-0841, 073-0731674-1673 and 079-0790462-0450).

#### Appendix A. Supplementary data

Supplementary data associated with this article can be found, in the online version, at <http://dx.doi.org/10.1016/j.aquatox.2016.03.009>.

#### References

- ATSDR, 2004. *Toxicological profile for cobalt*. U.S. Department of Health and Human Services, Atlanta.
- Appenroth, K.-J., Stöckel, J., Srivastava, A., Strasser, R., 2001. Multiple effects of chromate on the photosynthetic apparatus of *Spirodela polyrrhiza* as probed by OJIP chlorophyll a fluorescence measurements. *Environ. Pollut.* 115, 49–64, [http://dx.doi.org/10.1016/S0269-7491\(01\)00091-4](http://dx.doi.org/10.1016/S0269-7491(01)00091-4).
- Assche, F., Clijsters, H., 1990. Effects of metals on enzyme activity in plants. *Plant Cell Environ.* 13, 195–206, <http://dx.doi.org/10.1111/j.1365-3040.1990.tb01304.x>.
- Babić, M., Radić, S., Cvjetko, P., Roje, V., Pevalak-Kozlina, B., Pavlica, M., 2009. Antioxidative response of *Lemna minor* plants exposed to thallium (I)-acetate. *Aquat. Bot.* 91, 166–172, <http://dx.doi.org/10.1016/j.aquabot.2009.05.005>.
- Bakkaus, E., Gouget, B., Gallien, J.-P., Khodja, H., Carrot, F., Morel, J., Collins, R., 2005. Concentration and distribution of cobalt in higher plants: The use of micro-PIXE spectroscopy. *Nucl. Instrum. Meth. B* 231, 350–356, <http://dx.doi.org/10.1016/j.nimb.2005.01.082>.
- Belatik, A., Hotchandani, S., Carpentier, R., 2013. A. Belatik S. Hotchandani R. Carpentier Inhibition of the water oxidizing complex of photosystem II and the reoxidation of the quinone acceptor Q<sub>A</sub><sup>-</sup> by Pb<sup>2+</sup>. *PLoS One*, e68142, <http://dx.doi.org/10.1371/journal.pone.0068142>.
- Bradford, M.M., 1976. A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein–dye binding. *Anal. Biochem.* 72, 248–254, [http://dx.doi.org/10.1016/0003-2697\(76\)90527-3](http://dx.doi.org/10.1016/0003-2697(76)90527-3).
- Brestic, M., Zivcak, M., 2013. PSII fluorescence techniques for measurement of drought and high temperature stress signal in plants: protocols and applications. In: Rout, G.R., Das, A.B. (Eds.), *Molecular Stress Physiology of Plants*. Springer, Dordrecht, pp. 87–131 [http://dx.doi.org/10.1007/978-81-322-0807-5\\_4](http://dx.doi.org/10.1007/978-81-322-0807-5_4).
- Chatterjee, J., Chatterjee, C., 2000. Phytotoxicity of cobalt, chromium and copper in cauliflower. *Environ. Pollut.* 109, 69–74, [http://dx.doi.org/10.1016/S0269-7491\(99\)00238-9](http://dx.doi.org/10.1016/S0269-7491(99)00238-9).
- Chatterjee, J., Chatterjee, C., 2003. Management of phytotoxicity of cobalt in tomato by chemical measures. *Plant Sci.* 164, 793–801, [http://dx.doi.org/10.1016/S0168-9452\(03\)00066-9](http://dx.doi.org/10.1016/S0168-9452(03)00066-9).
- Chen, L., Jia, H., Tian, Q., Du, L., Gao, Y., Miao, X., Liu, Y., 2012. Protecting effect of phosphorylation on oxidative damage of D1 protein by down-regulating the production of superoxide anion in photosystem II membranes under high light. *Photosynth. Res.* 112, 141–148, <http://dx.doi.org/10.1007/s11120-012-9750-9>.
- Csatorday, K., Gombos, Z., Szalontai, B., 1984. Mn<sup>2+</sup> and Co<sup>2+</sup> toxicity in chlorophyll biosynthesis. *Proc. Natl. Acad. Sci. U. S. A.* 81, 476–478.
- De Ronde, J., Cress, W., Krüger, G., Strasser, R., Van Staden, J., 2004. Photosynthetic response of transgenic soybean plants, containing an *Arabidopsis P5CR* gene, during heat and drought stress. *J. Plant Physiol.* 161, 1211–1224, <http://dx.doi.org/10.1016/j.jplph.2004.01.014>.
- Drzewiecka-Matuszek, A., Skalna, A., Karocki, A., Stochel, G., Fiedor, L., 2005. Effects of heavy central metal on the ground and excited states of chlorophyll. *J. Biol. Inorg. Chem.* 10, 453–462, <http://dx.doi.org/10.1007/s00775-005-0652-6>.
- EBRC, ARCHE, 2012. *Exposure Scenarios Cobalt Update*. EBRC – ARCHE, Hannover-Gent.
- El-Sheekh, M.M., El-Naggar, A.H., Osman, M.E.H., El-Mazaly, E., 2003. Effect of cobalt on growth, pigments and the photosynthetic electron transport in *Monoraphidium minutum* and *Nitzschia perminuta*. *Braz. J. Plant Physiol.* 15, 159–166, <http://dx.doi.org/10.1590/S1677-04202003000300005>.
- Falqueto, A.R., dos Santos, P.N., Fontes, R.V., Silva, D.M., 2013. Analysis of chlorophyll a fluorescence of two mangrove species of Vitória Bay (ES, Brazil) to natural variation of tide. *Revista Biociências*, 18.
- Gad, N., Hassan, N.M., 2013. Response of growth and yield of sweet pepper (*Capiscum annuum* L.) to cobalt nutrition. *World Appl. Sci. J.* 21, 760–765, <http://dx.doi.org/10.5829/idosi.wasj.2013.21.5.2890>.
- Giannopolitis, C.N., Ries, S.K., 1977. Superoxide dismutases I. Occurrence in higher plants. *Plant Physiol.* 59, 309–314.
- Gopal, R., 2014. Antioxidant defense mechanism in pigeon pea under cobalt stress. *J. Plant Nutr.* 37, 136–145, <http://dx.doi.org/10.1080/01904167.2013.849734>.

- Hasni, I., Hamdani, S., Carpentier, R., 2013. Destabilization of the oxygen evolving complex of photosystem II by  $Al^{3+}$ . *Photochem. Photobiol.* 89, 1135–1142 <http://dx.doi.org/10.1111/php.12116>.
- Horvatić, J., Peršić, V., 2007. The effect of  $Ni^{2+}$ ,  $Co^{2+}$ ,  $Zn^{2+}$ ,  $Cd^{2+}$  and  $Hg^{2+}$  on the growth rate of marine diatom *Phaeodactylum tricornutum* Bohlin: microplate growth inhibition test. *Bull. Environ. Contam. Toxicol.* 79, 494–498. <http://dx.doi.org/10.1007/s00128-007-9291-7>.
- Ince, N.H., Dirilgen, N., Apikyan, I.G., Tezcanli, G., Üstün, B., 1999. Assessment of toxic interactions of heavy metals in binary mixtures: a statistical approach. *Arch. Environ. Contam. Toxicol.* 36, 365–372 <http://dx.doi.org/10.1007/PL00006607>.
- Joshi, M.K., Mohanty, P., 2004. Chlorophyll *a*. In: Papageorgiou, G.C., Govinjee (Eds.), *Chlorophyll a fluorescence: A signature of photosynthesis*. Springer, Dordrecht, pp. 637–661 [http://dx.doi.org/10.1007/978-1-4020-3218-9\\_25](http://dx.doi.org/10.1007/978-1-4020-3218-9_25).
- Khan, M.R., Khan, M.M., 2010. Effect of varying concentration of nickel and cobalt on the plant growth and yield of chickpea. *Aust. J. Basic Appl. Sci.* 4, 1036–1046.
- Lazár, D., Ilík, P., Nauš, J., 1997. An appearance of K-peak in fluorescence induction depends on the acclimation of barley leaves to higher temperatures. *J. Lumin.* 72–74, 595–596 [http://dx.doi.org/10.1016/S0022-2313\(96\)00293-1](http://dx.doi.org/10.1016/S0022-2313(96)00293-1).
- Lepeduš, H., Gača, V., Viljevac, M., Kovač, S., Fulgosi, H., Šimić, D., Jurković, V., Cesar, V., 2011. H. Lepeduš V. Gača M. Viljevac S. Kovač H. Fulgosi D. Šimić V. Jurković V. Cesar Changes in photosynthetic performance and antioxidative strategies during maturation of Norway maple (*Acer platanoides* L.) leaves. *Plant Physiol. Bioch.* 49, 368–376. <http://dx.doi.org/10.1016/j.plaphy.2010.12.011>.
- Lichtenthaler, H.K., 1987. Chlorophylls and carotenoids: pigments of photosynthetic biomembranes. *Methods Enzymol.* 148C, 350–382. [http://dx.doi.org/10.1016/0076-6879\(87\)48036-1](http://dx.doi.org/10.1016/0076-6879(87)48036-1).
- Lichtenthaler, H.K., 1996. Vegetation stress: an introduction to the stress concept in plants. *J. Plant Physiol.* 148, 4–14. [http://dx.doi.org/10.1016/S0176-1617\(96\)80287-2](http://dx.doi.org/10.1016/S0176-1617(96)80287-2).
- Mallick, N., Mohn, F., 2003. N. Mallick F. Mohn Use of chlorophyll fluorescence in metal-stress research: a case study with the green microalga *Scenedesmus*. *Ecotox. Environ. Saf.* 55, 64–69. [http://dx.doi.org/10.1016/S0147-6513\(02\)00122-7](http://dx.doi.org/10.1016/S0147-6513(02)00122-7).
- Milone, M.T., Sgherri, C., Clijsters, H., Navari-Izzo, F., 2003. Antioxidative responses of wheat treated with realistic concentration of cadmium. *Environ. Exp. Bot.* 50, 265–276. [http://dx.doi.org/10.1016/S0098-8472\(03\)00037-6](http://dx.doi.org/10.1016/S0098-8472(03)00037-6).
- Mohanty, N., Vass, I., Demeter, S., 1989. Impairment of photosystem 2 activity at the level of secondary quinone electron acceptor in chloroplasts treated with cobalt, nickel and zinc ions. *Physiol. Plant.* 76, 386–390. <http://dx.doi.org/10.1111/j.1399-3054.1989.tb06208.x>.
- Nagajyoti, P.C., Lee, K.D., Sreekanth, T.V.M., 2010. Heavy metals, occurrence and toxicity for plants: a review. *Environ. Chem. Lett.* 8, 199–216. <http://dx.doi.org/10.1007/s10311-010-0297-8>.
- Naumann, B., Eberius, M., Appenroth, K.J., 2007. Growth rate based dose-response relationships and EC-values of ten heavy metals using the duckweed growth inhibition test (ISO 20079) with *Lemna minor* L. clone St. *J. Plant Physiol.* 164, 1656–1664. <http://dx.doi.org/10.1016/j.jplph.2006.10.011>.
- Osman, M.E.H., El-Naggar, A.H., El-Sheekh, M.M., El-Mazally, E.E., 2004. Differential effects of  $Co^{2+}$  and  $Ni^{2+}$  on protein metabolism in *Scenedesmus obliquus* and *Nitzschia perminuta*. *Environ. Toxicol. Pharmacol.* 16, 169–178. <http://dx.doi.org/10.1016/j.etap.2003.12.004>.
- Oukarroum, A., Madidi, S.E., Schansker, G., Strasser, R.J., 2007. Probing the responses of barley cultivars (*Hordeum vulgare* L.) by chlorophyll *a* fluorescence OLKJIP under drought stress and re-watering. *Environ. Exp. Bot.* 60, 438–446. <http://dx.doi.org/10.1016/j.envexpbot.2007.01.002>.
- Outridge, P., Noller, B., 1991. Accumulation of toxic trace elements by freshwater vascular plants. In: Ware, G.W. (Ed.), *Reviews of Environmental Contamination and Toxicology*. Springer, New York, pp. 1–63 [http://dx.doi.org/10.1007/978-1-4612-3196-7\\_1](http://dx.doi.org/10.1007/978-1-4612-3196-7_1).
- Palit, S., Sharma, A., Talukder, G., 1994. Effects of cobalt on plants. *Bot. Rev.* 60, 149–181. <http://dx.doi.org/10.1007/BF02856575>.
- Pan, X., Zhang, D., Chen, X., Li, L., Mu, G., Li, L., Song, W., 2010. Sb uptake and photosynthesis of *Zea mays* growing in soil watered with Sb mine drainage: an OJIP chlorophyll fluorescence study. *Pol. J. Environ. Stud.* 19, 981–987.
- Panda, S.K., 2008. Impact of copper on reactive oxygen species, lipid peroxidation and antioxidants in *Lemna minor*. *Biol. Plantarum* 52, 561–564. <http://dx.doi.org/10.1007/s10535-008-0111-7>.
- Perreault, F., Popovic, R., Dewez, D., 2014. Different toxicity mechanisms between bare and polymer-coated copper oxide nanoparticles in *Lemna gibba*. *Environ. Pollut.* 185, 219–227. <http://dx.doi.org/10.1016/j.envpol.2013.10.027>.
- Pirson, A., Seidel, F., 1950. Zell- und stoffwechselfysiologische Untersuchungen an der Wurzel von *Lemna minor* L. unter besonderer Berücksichtigung von Kalium- und Kalziummangel. *Planta* 38, 431–473. <http://dx.doi.org/10.1007/BF01928941>.
- Plekhanov, S., Chemeris, Y.K., 2003. Early toxic effects of zinc, cobalt, and cadmium on photosynthetic activity of the green alga *Chlorella pyrenoidosa* Chick S-39. *Biol. Bull. Russian Acad. Sci.* 30, 506–511. <http://dx.doi.org/10.1023/A.1025806921291>.
- Pospíšil, P., 2012. Molecular mechanisms of production and scavenging of reactive oxygen species by photosystem II. *Biochim. Biophys. Acta-Bioenerg.* 1817, 218–231. <http://dx.doi.org/10.1016/j.bbabi.2011.05.017>.
- Prasad, M.N.V., Malec, P., Waloszek, A., Bojko, M., Strzalka, K., 2001. Physiological responses of *Lemna trisulca* L. (duckweed) to cadmium and copper bioaccumulation. *Plant Sci.* 161, 881–889. [http://dx.doi.org/10.1016/S0168-9452\(01\)00478-2](http://dx.doi.org/10.1016/S0168-9452(01)00478-2).
- Prasad, M.N.V., 2004. *Heavy Metal Stress in Plants: From Biomolecules to Ecosystems*. Springer Berlin Heidelberg.
- Razinger, J., Dermastia, M., Koce, J.D., Zrimec, A., 2008. Oxidative stress in duckweed (*Lemna minor* L.) caused by short-term cadmium exposure. *Environ. Pollut.* 153, 687–694. <http://dx.doi.org/10.1016/j.envpol.2007.08.018>.
- Sandalio, L., Dalurzo, H., Gomez, M., Romero-Puertas, M., Del Rio, L., 2001. Cadmium-induced changes in the growth and oxidative metabolism of pea plants. *J. Exp. Bot.* 52, 2115–2126. <http://dx.doi.org/10.1093/jxbbot/52.364.2115>.
- Shalygo, N., Kolesnikova, N., Voronetskaya, V., Averina, N., 1999. Effects of  $Mn^{2+}$ ,  $Fe^{2+}$ ,  $Co^{2+}$ , and  $Ni^{2+}$  on chlorophyll accumulation and early stages of chlorophyll formation in greening barley seedlings. *Russ. J. Plant Physiol.* 46, 496–501.
- Sree, K.S., Keresztes, Á., Mueller-Roeber, B., Brandt, R., Eberius, M., Fischer, W., Appenroth, K.-J., 2015. Phytotoxicity of cobalt ions on the duckweed *Lemna minor* – Morphology, ion uptake, and starch accumulation. *Chemosphere* 131, 149–156. <http://dx.doi.org/10.1016/j.chemosphere.2015.03.008>.
- Srivastava, A., Guissé, B., Greppin, H., Strasser, R.J., 1997. Regulation of antenna structure and electron transport in photosystem II of *Pisum sativum* under elevated temperature probed by the fast polyphasic chlorophyll *a* fluorescence transient: OKJIP. *Biochim. Biophys. Acta-Bioenerg.* 1320, 95–106. [http://dx.doi.org/10.1016/S0005-2728\(97\)00017-0](http://dx.doi.org/10.1016/S0005-2728(97)00017-0).
- Strasser, R.J., Srivastava, A., Tsimilli-Michael, M., 2000. The fluorescence transient as a tool to characterize and screen photosynthetic samples. In: Yunus, M., Pathre, U., Mohanty, P. (Eds.), *Probing Photosynthesis: Mechanism, Regulation & Adaptation*. Probing photosynthesis: Mechanism, regulation & adaptation. 1st ed. CRC, New York, pp. 445–483.
- Strasser, R.J., Tsimilli-Michael, M., Srivastava, A., 2004. Analysis of the chlorophyll *a* fluorescence transient. In: Papageorgiou, G.C., Govinjee (Eds.), *Chlorophyll a Fluorescence: A signature of photosynthesis*. Springer, Dordrecht, pp. 321–362. [http://dx.doi.org/10.1007/978-1-4020-3218-9\\_12](http://dx.doi.org/10.1007/978-1-4020-3218-9_12).
- Tóth, S.Z., Schansker, G., Garab, G., Strasser, R.J., 2007. Photosynthetic electron transport activity in heat-treated barley leaves: the role of internal alternative electron donors to photosystem II. *Biochim. Biophys. Acta-Bioenerg.* 1767, 295–305. <http://dx.doi.org/10.1016/j.bbabi.2007.02.019>.
- Tewari, R.K., Kumar, P., Sharma, P.N., Bisht, S.S., 2002. Modulation of oxidative stress responsive enzymes by excess cobalt. *Plant Sci.* 162, 381–388. [http://dx.doi.org/10.1016/S0168-9452\(01\)00578-7](http://dx.doi.org/10.1016/S0168-9452(01)00578-7).
- Tkalec, M., Prebeg, T., Roje, V., Pevalek-Kozlina, B., Ljubesic, N., 2008. Cadmium-induced responses in duckweed *Lemna minor* L. *Acta Physiol. Plant.* 30, 881–890. <http://dx.doi.org/10.1007/s11738-008-0194-y>.
- Velikova, V., Tsonev, T., Loreto, F., Centritto, M., 2011. Changes in photosynthesis, mesophyll conductance to  $CO_2$ , and isoprenoid emissions in *Populus nigra* plants exposed to excess nickel. *Environ. Pollut.* 159, 1058–1066. <http://dx.doi.org/10.1016/j.envpol.2010.10.032>.
- Verma, S., Dubey, R., 2003. Lead toxicity induces lipid peroxidation and alters the activities of antioxidant enzymes in growing rice plants. *Plant Sci.* 164, 645–655. [http://dx.doi.org/10.1016/S0168-9452\(03\)00022-0](http://dx.doi.org/10.1016/S0168-9452(03)00022-0).
- Vidaković-Cifrek, Ž., Sorić, S., Babić, M., 2013. Growth and photosynthesis of *Lemna minor* L. exposed to different light conditions and sucrose supplies. *Acta Bot. Croat.* 72, 211–219. <http://dx.doi.org/10.2478/v10184-012-0018-4>.
- Vidaković-Cifrek, Ž., Tkalec, M., Šikić, S., Tolić, S., Lepeduš, H., Pevalek-Kozlina, B., 2015. Growth and photosynthetic responses of *Lemna minor* exposed to cadmium in combination with zinc or copper. *Arch. Indu. Hygiene Toxicol.* 66. <http://dx.doi.org/10.1515/aiht-2015-66-2618>.
- Wojtaszek, P., 1997. Oxidative burst: an early plant response to pathogen infection. *Biochem. J.* 322, 681–692. <http://dx.doi.org/10.1042/bj3220681>.
- Yusuf, M.A., Kumar, D., Rajwanshi, R., Strasser, R.J., Tsimilli-Michael, M., Govindjee, Sarin, N.B., 2010. Overexpression of gamma-tocopherol methyl transferase gene in transgenic *Brassica juncea* plants alleviates abiotic stress: physiological and chlorophyll *a* fluorescence measurements. *Biochim. Biophys. Acta*, 1428–1438. <http://dx.doi.org/10.1016/j.bbabi.2010.02.002>.
- Zeid, I.M., Ghazi, S., Nabawy, D., 2013. Alleviation of Co and Cr toxic effects on alfalfa. *Int. J. Agron. Plant Product.* 4, 984–993.
- Zeid, I., 2001. Responses of *Phaseolus vulgaris* chromium and cobalt treatments. *Biol. Plantarum* 44, 111–115. <http://dx.doi.org/10.1023/A:1017934708402>.